An experimental framework for ecosystem capital accounting in Europe

ISSN 1725-2237

An experimental framework for ecosystem capital accounting in Europe

Cover design: EEA

Layout: EEA/Henriette Nilsson

Legal notice

The contents of this publication do not necessarily reflect the official opinions of the European Commission or other institutions of the European Union. Neither the European Environment Agency nor any person or company acting on behalf of the Agency is responsible for the use that may be made of the information contained in this report.

Copyright notice

© EEA, Copenhagen, 2011

Reproduction is authorised, provided the source is acknowledged, save where otherwise stated.

Information about the European Union is available on the Internet. It can be accessed through the Europa server (www.europa.eu).

Luxembourg: Publications Office of the European Union, 2011

ISBN 978-92-9213-233-0 ISSN 1725-2237 doi:10.2800/83749

By Jean-Louis Weber

Special Adviser on Economic-Environmental Accounting European Environment Agency



European Environment Agency Kongens Nytorv 6 1050 Copenhagen K Denmark

Tel.: +45 33 36 71 00 Fax: +45 33 36 71 99 Web: eea.europa.eu

Enquiries: eea.europa.eu/enquiries

Contents

For	ew	ord	6
Exe	cut	tive summary	7
	1.1	E purpose of environmental-economic accounting Environmental-economic accounts and national accounts Simplified ecosystem capital accounts	8
	2.1 2.2 2.3	Statistical units and classifications	12 17 17
		graphy 1 List of tables and accounting items	
		2 Mock-up accounts	

Foreword

The need to account for natural resources as capital, in the same way as we account for economic and financial resources, is getting more and more attention. The recently published EU Roadmap for a Resource Efficient Europe sets the policy framework for action in the coming years and decades for which robust data and indicators are needed.

In this respect, there is a contrast between the natural resources which are recorded by the System of National Accounts 2008, the basis for GDP, and other natural resources which are ignored because they are not seen as economic assets by the market. The latter is typically the case for many important ecosystem functions and services which contribute to people's wellbeing but are considered as external by the economy.

To bridge this gap, the UN Committee of Experts on Environmental-Economic Accounting decided in its June 2011 meeting to include experimental ecosystem accounts in the scope of the revision of the System of Environmental-Economic Accounts revision by 2013. In Europe, a project to test the feasibility of ecosystem capital accounts was launched by the EEA in 2010 in anticipation of such stakeholder demands.

The experimental accounts framework presented in this report builds upon the experience gained in this feasibility study. It has also benefited from the reflections which have taken place in the UN SEEA revision process as well as from the European Union GDP and Beyond initiative and TEEB, The Economics of Ecosystems and Biodiversity, process. The EEA has been actively participating in these activities and has inter-alia, produced a report on ecosystem accounting for coastal Mediterranean wetlands as well as contributed to the TEEB D1 report for National and International Policy Making.

The ecosystem accounting framework presented here is being tested in the context of an open Europe taking stock of its relations to the rest of the world. Because ecosystem accounts are deep-rooted into monitoring databases, implementation presently focuses on physical accounts. Monetary valuations, the adjustment of national accounts aggregates for income and final consumption, and the calculation of ecological debts, are foreseen in subsequent steps within the same logical framework. Although designed for European needs, most of the features proposed here have a global scope, and so can support experimentation in other regions of the world.

The progress made so far has been possible through the support of EEA partners and networks and the developments of environmental information systems in recent years. Ecosystem accounting is a module under the European Strategy for Environmental Accounting managed by Eurostat. The basis for physical accounts for Europe is the Corine land cover inventory 1990–2006 produced in collaboration with Eionet, the European Space Agency and the Joint Research Centre. EEA's own databases have been mined for computing the test accounts alongside the databases of Eurostat and the Joint Research Centre.

The EEA Management Board, Eionet and the EEA Scientific Committee have been consulted regularly and provided guidance. Last but not least, the EEA policy framework has constantly confronted stated policy requests regarding national accounts, efficient resource use or nature conservation and methodological proposals. I would like to thank all for their contributions and look forward to continuing cooperation in coming years.

Prof. Jacqueline McGlade

Executive Director European Environment Agency

Executive summary

Ecosystem accounts are being developed as part of the System of Environmental-Economic Accounts which aims at supplementing the UN System of National Accounts with information on the environment and natural capital. The purpose is to broaden the scope of the variables taken into account in policymaking in order to improve understanding of the interdependence and interactions between the economy and the environment. Ultimately, these ecosystem accounts will yield new indicators and aggregates expressed in physical and monetary units that will be made available to policymakers and analysts to assess the efficiency of natural resource use, the pattern of economic growth, the contribution of nature and its use within and outside the market, the short- and longer-term constraints resulting from the need to maintain living and other renewable capital, and the related benefits and costs.

At the end of 2009, the European Environment Agency launched an experimental project to implement simplified ecosystem capital accounts for Europe as a 'fast-track' initiative, based on the use of existing data and statistics. In addition to feasibility assessment, the project aims at framing ecosystem accounts and identifying which indicators and aggregates could be delivered and integrated into enlarged national accounts. Based on the project findings, an overall framework for ecosystem capital accounting has been designed. It highlights accounting balances and relationships between accounting tables and systems as well as key indicators and aggregates that describe economy-ecosystem interactions.

The indicators and aggregates include: the **ecosystem resource accessible surplus** (which

shows the level of resources that can be used without jeopardising ecosystem reproduction functions); the demand for (accessible) ecosystem services per capita, which is a measure of ecosystem contribution to well-being; the total ecosystem capital potential, defined as the biomass accessible under the constraints of maintaining accessibility to water, green landscape infrastructure and biodiversity (and measured in a 'numeraire' referred to as the Ecosystem Potential Unit Equivalent); the Ecosystem Capital Degradation (ECD) which describes domestic ecosystem overuse; Consumption of Ecosystem Capital (CEC, the ecosystem capital depreciation in SNA terminology), calculated as (physical) ECD valued by remediation costs; and the equivalent ECD embedded in imports and exports for commodities produced in unsustainable conditions. As a next stage it is proposed to use CEC to adjust National Accounts aggregates: CEC Adjusted Net Domestic Product or CEC Adjusted Net National Income, Final Consumption at Full Cost (including non-paid CEC), Imports and Exports at Full Cost. Using this approach, two balance sheets of assets and liabilities are ultimately established, one in physical units, the other in terms of money. The balance sheets of financial liabilities allow, amongst other things, a record to be kept of the amount of ecological debt first in physical units regarding physical degradation and second in monetary units to balance the non-paid consumption of ecosystem capital. Last but not least, recording ecological debts makes it possible to keep the conventional GDP unchanged while supplementing it with appropriate adjusted aggregates.

1 The purpose of environmentaleconomic accounting

1.1 Environmental-economic accounts and national accounts

The purpose of environmental-economic accounting is to supplement the conventional national accounts (UN SNA 2008) with tables which inform policymakers of environmental and natural resource availability, use, depletion and degradation. Through such accounts, economic performance measured by aggregates like Gross Domestic Product, Net National Income, Final Consumption, Net Savings, Imports and Exports, Assets and Liabilities or Employment can be balanced by natural capital indicators that describe the opportunities and constraints, benefits and costs, efficiency of resource use, and externalities that arise in relation to interactions with the environment. Implementation of environmental-economic accounts has been recognised as an important step towards: sustainable development (see Agenda 21 of 1992); the measurement of economic progress (see Beyond GDP and the so-called Stiglitz-Sen-Fitoussi commission); support for green economy (UNEP) and green growth (OECD) strategies; designing resource efficiency policies (see UNEP and EC's Flagship Initiative for a resource-efficient Europe); and for biodiversity conservation (see the Aichi-Nagoya CBD's strategy of 2010). In 2007, the UN Statistical Commission mandated the UN Committee of Experts on Environmental-Economic Accounts (UNCEEA) to raise the 'Integrated System of Environmental-Economic Accounts' (SEEA 2003) up to the level of an international standard by 2012.

Although environmental-economic accounts are compiled using both physical and monetary units, the former are considered the basis of the framework in the EEA initiative.

Accounts in physical units aim primarily at supplementing conventional national accounts with data on the use and availability of natural resources. The objective is to measure the overall efficiency of the economy, first in terms of the material or energy resource input (and waste generation) necessary to produce one unit of GDP, and second to assess resource depletion. Using this approach, physical constraints can be incorporated into macro-economic analysis and support action

towards **greener** growth, development and actions, both public (e.g. via taxes, regulation and planning) and private (e.g. via productivity gains, technology, contents of final consumption).

Physical units can be specific to the kind of resource recorded (tonnes, cubic meters, hectares, number of units) or common to a range of resources. In this case, a unit-equivalent needs to be found. For example, material flow accounts which are currently the main basis for resource-efficiency analysis record 'everything' in tonnes. Another solution is to use carbon or energy unit-equivalents, as in UNFCCC reporting. The Ecological Footprint Accounts propose surface area as a general unit-equivalent. These solutions are obviously incomplete (e.g. economy-wide material flow accounts commonly set aside water, considered as 'so large that they would dominate all other materials' (1), and land, which has no mass in itself) and limited in scope due to the specific equivalence functions used. Moreover, the qualitative aspects of the living natural resource are broadly ignored, nature being considered only as a 'mine' for resources, and so subject to depletion rather than degradation of its capability for self-renewal.

Monetary values does not ignore the qualities of assets and commodities. However, not all qualities are considered, only those which matter to economic actors in their search of profit and wealth. In the case of subsoil assets, the issue can be neglected as long they exist only as an economic resource. Considering assets with multiple functions like dynamic, biophysical systems, which can be regarded as an economic resource and a public good, market valuation does not encompass all the elements of present and future scarcity needed to assess green growth options and so frame green economy scenarios. Generally, one 'main' function is considered as productive and used to capture most of (if not all) the economic value; other functions being considered as free externalities or ignored. In the absence of external enforcement of such values (e.g. via environmental taxes or norms) market prices are incomplete and beyond market values although they are part of human wellbeing and should be included in any assessment of sustainable development.

 $^{(^1) \}quad \text{See http://epp.eurostat.ec.europa.eu/statistics_explained/index.php/Material_flow_accounts.}$

The extension of national accounts to cover economic natural assets and their services (incorporated into commodities) is important but cannot deliver a sufficiently complete vision of the interaction of people and nature. For example, an enterprise holding and managing a forest will know and care about trees and timber, but much less about 'non-timber forest values', or forest water regulating functions and micro-climate effects which may be highly important for other sectors of society and for biodiversity. One reason is that the forest is privately managed for private benefits, while the other ecosystem services are mostly public goods. Another reason is that large parts of nature are out of the scope of those 'owned and managed for profit' which is the category of natural assets recorded in the SNA. For these reasons, the UNCEEA decided in June 2011 to devote volume 2 of the new SEEA to ecosystem accounts. The volume will include accounts of flows and stocks in physical units, and where relevant and consistent with SNA principles, valuations. The EEA, Eurostat and the World Bank have been asked to support the preparation of SEEA volume 2. This paper reports progress in Europe on ecosystem capital accounts, which express the capability of ecosystems to contribute alongside other forms of capital and to deliver services, and the responsibility of the economy regarding their good maintenance.

1.2 Simplified ecosystem capital accounts

The purpose of developing ecosystem capital accounts is to assess the sustainability of the economy-ecosystem interaction from the standpoint of nature, to measure the state of the ecosystems, and, when degradation is observed, to calculate the costs of avoiding damage, or of repair and compensation. These can all be regarded as measurements of ecosystem capital depreciation or 'consumption' (in the SNA sense). In such a setting, physical accounts provide a measure of the physical constraints that cannot be surpassed by the economy without causing damage to human communities and the economy itself.

At the end of 2009, the EEA launched an experimental project 'fast-track implementation of simplified ecosystem capital accounts' for Europe — 'fast-track' because of urgent and recurrent policy demands and 'simplified', because full details are not necessary at the macro level. The approach adopted is top-down, based on Europe-wide datasets and statistics but, as far as possible, data and statistics are compiled at the level of the standard European

1km² grid. The use of the grid is justified by requirements of change detection, and the flexibility needed to report in terms of different geographical units (e.g. regions, river basins, coastal zones). The approach also anticipates the forthcoming expected links with accounting applications at the national level. The test is carried out with existing data and statistics, with the aim of supplying annual updates (to meet the policy agenda) and retrospective time series. Physical accounts are being developed and computed first; the valuation of costs and benefits is still at an exploratory stage. The framework developed for SECA in Europe is an input to the current preparation of SEEA volume 2.

The narrative behind Simplified Ecosystem Capital Accounts (SECA): Ecosystems can be described as capital which delivers a bundle of services to people, some of which are appropriated and incorporated into products, accumulated and/or consumed. Other services are public goods of common benefit to the economy and human wellbeing. Altogether, these ecosystem services depend on ecosystem capital regeneration which is in turn influenced by ecosystem services consumption.

In the fact-track accounts three groups of ecosystem services have been considered: accessible biomass/ carbon, accessible water, and accessible regulating and cultural services. Accessible refers to the share of the 'total' or 'available' resource which can be used without damaging ecosystem capital capacity. All three groups of services are generally produced (in variable proportions) by all ecosystems. Accessible biomass/carbon and water together make up 99 per cent of all 'provisioning services' as described in the Millennium Ecosystem Assessment (MA) or Common International Classification of Ecosystem Services (CICES) classifications of ecosystem services. Biomass/carbon and water are recorded in formal balances while regulating and cultural services are measured indirectly on the basis of ecosystem capacity to deliver them (state of landscape green infrastructure and biodiversity). For each of these groups, the amount of services which can be used must be lower than the accessible surplus, which means that in terms of sustainable development there should not be significant trade-offs between these services.

The primary ecosystem service is production of biomass which can be generated and withdrawn (by agriculture, forestry, fisheries, etc.) up to a surplus which takes into account nature's own reproductive needs. The surplus corresponds to the current 'food of biodiversity' and the maintenance of bio-carbon stocks in soil and perennial vegetation,

and which is required if the ecosystem is to be self-sustaining. Production of biomass must also be compatible with the maintenance of accessible water resources (e.g. limits to irrigation) and the bundle of services supplied by the green landscape infrastructure. Similarly, water can be abstracted only up to an accessible surplus, to ensure the good functioning of the water cycle, as well as biomass production, and the needs of landscapes and biodiversity; for example, a new reservoir destroys previous ecosystem functions, over-dimensioned irrigation infrastructures create risks of agricultural shortages in years with rainfall deficit. The development of landscape services may result in the reduction, for example, of biomass production because of subsequent falling yields — which will be recorded in the carbon/biomass account.

'Accessible' means that not all the available resource can be used because of physical constraints (a large part of the aquifers, flood water greater than needed for reservoirs replenishment), inappropriate location or timeliness, inappropriate quality, and because part of the annual service flow has to be left to the ecosystem for its own needs. In the case of services supplied by green landscape infrastructure, accessibility is dependent on the population which can access it and the inverse of landscape artificiality (including urban areas where most of the population on Europe lives). The concept of resource accessibility is particularly important regarding the demand for ecosystem services and the definition of robust indicators with clear definitions of the limits of sustainable use. Such indicators implemented at the appropriate scale can be associated with population data, considerably increasing their usefulness for policymaking.

The calculation of **Total Ecosystem Potential**, **Net Change in TEP and Ecosystem Capital Degradation** summarises the state of the ecosystem capital. Total inland, sea and atmosphere ecosystem potential measured in the basic balance in tonnes of carbon is weighted by a set of composite indexes which reflect the external factors that limit carbon accessibility: Ecosystem Accessible Water Surplus (EAWS), Landscape Ecosystem Potential (LEP), River Ecosystem Potential (2) (REP) and Ecosystem Biodiversity Rating (EBR) (which combines green infrastructure and species diversity measurements). It results in a new unit called **EPUE** for **Ecosystem Potential Unit Equivalent**. Gain in EPUE means

positive effects of restoration programmes and/ or natural improvement, loss means degradation by activities and/or natural disturbance. Particular attention is given to the calculation of **Ecosystem** Capital Degradation (ECD) which is the result of economic activity and will be used in a subsequent step to calculate ecosystem capital depreciation. Ecosystem capital degradation is for that purpose analysed in a special table according to the stress factors that have caused it: land-cover change, restructuring/de-structuring of landscapes and rivers, over-exploitation of biological resources, waste disposal, and pollution. It is then possible to calculate, factor by factor, the cost of restoring one unit of EPUE. Depending on the ecosystems and issues being considered, costs will reflect reductions in yields, abatement of pollution (including GHGs), and programmes such as the replanting of hedgerows and reforestation. In the accounts, cost calculations are based on observed practices not on individual preferences.

Back to GDP

Consumption of Ecosystem Capital is similar to Consumption of Fixed Capital (CFC) and should be treated in a similar way as a deduction when shifting from Gross Domestic Product to Net Domestic Product or Net National Income. Another approach is to consider that, unlike CFC, which is included in the value of economic assets and therefore transferred to the value of commodities, CEC is not paid. This means that CEC is not included in the purchaser price of Final Consumption, nor in Imports and Exports. This major price distortion can be corrected by adding up the unpaid CECs to calculate Final Consumption, Imports and Exports at Full Cost. This would not require changing the conventional calculation of GDP, the CEC price adjustment being balanced by an appropriate recording of ecological debts.

The proposed way of calculating ecosystem capital degradation (or CEC) diverges from dominant economic theory which defines depreciation as a loss in asset value which is equivalent (in the absence of reliable market prices for assets, which is generally the case for natural capital) to the discounted net expected future benefits (net present value). The difficulty of the latter method at the macro scale is that it implies assessment

⁽²⁾ In the case of rivers, the potential is equivalent to exergy (or accessible energy regarding water position or potential, its concentration in various substances and other elements such as temperature or speed — see Valero, 2006), measured in energy units.

and valuation of all individual services provided by the multiple functions of ecosystems and their aggregation without double counting. There is no evidence so far that this conventional method, implemented successfully in many case studies, can be used for national accounting. The proposed approach, which combines physical degradation and restoration costs, is probably just a surrogate for the one prescribed by economic theory, but its implementation seems feasible.

2 The simplified framework of ecosystem capital accounts

Experimental implementation of SECA in Europe and the preliminary discussions on ecosystem accounting at the international level have clarified the design of a simplified framework of ecosystem capital accounts. It has involved defining measurement and statistical units (3), classifications and an accounting structure.

2.1 Statistical units and classifications

Unlike analytical research and modelling, accounting requires crisp units with clear borders and stable classifications for compiling data and statistics and supporting comparisons in space and time (time series). The definition of such units and their classification is an essential preliminary step when defining an accounting framework.

2.1.1 Statistical units

The SNA defines basic units as legal entities entitled with complete capacity of taking any economic decision regarding production, consumption, investment, acquisition of financial assets or liabilities, etc. These 'institutional units' are typically enterprises, central or local government institutions or households. Regarding production analysis, the SNA chooses smaller units which are better correlated to particular products or groups of products, or more homogenous. They are called 'establishments'. An establishment is a part of an enterprise that is situated in a single location and which engages predominantly in one kind of economic activity.

Equivalent units need to be defined for ecosystems. Using specific statistical units for ecosystems instead of using economic or administrative units is a major step forward. In principle, ecosystems range from the microscopic level to the global. However, as ecosystem accounts are part of environmental-economic accounts, and aim at being used jointly with the SNA, priority should be given to equivalent levels for defining statistical units for ecosystems.

The scientific literature suggests that the best unit to assess ecosystems is the 'socio-ecological system (SES)' (Gallopin, 1991, Glaser, 2008). SES integrates ecosystem functions and dynamics as well as human activities and the interactions of all these. The SES is equivalent to the SNA's institutional unit. Considering the production of ecosystem services, and in particular provisioning services, SESs are more or less homogenous. A large forest is at the same time a socio-ecological system with its own behaviour and a unit of production of timber and most other ecosystem services. A small forest that is part of a mosaic landscape with agriculture, villages and natural areas is certainly a production unit for timber, but delivers other services only in conjunction with the neighbouring units; it is influenced by its environment and has less autonomy. Such units can be considered as equivalent to the establishments defined by the SNA.

Once this equivalence between SNA entities and SESs is accepted, the task is to define such entities in practice. SES and ecosystem production units are defined by their capacities to generate services, on a range of spatial scales. For the production of statistics, units need to have clear boundaries. There may be coincidences between the competency of institutional units and biophysical entities corresponding to one or other type foreseen. For example a natural reserve often covers an ecosystem, or a forest may belong to one single owner. But this is not the general case and another solution had to be found.

For inland ecosystems, the solution has been to analyse the biophysical characteristics of the landscapes. The production level can be addressed by mapping land-cover units. These are defined by their composition in terms of basic bio-physical objects or patches (e.g. grass, shrub, tree, rock and other minerals, sand, ice, snow and water), the type of use (artificial, cultivated, non-cultivated) and landscape patterns (fragmented, connected, etc.). The methodology has been developed by FAO under the name Land Cover Classification System (LCCS). In Europe, the Corine Land Cover classification

⁽³⁾ Measurement units are typically hectares, joules, cubic meters, 'ppm' or euros or \$. Statistical units are the entities for which statistics are collected — or accounts computed. They are typically enterprises and their establishments, government services or households.

follows similar principles and is being translated into the LCCS3 meta-language. One important point is that LCCS and Corine can be implemented using satellite images. The units for ecosystem accounting are named Land Cover Functional Units (LCFU).

Socio-Ecological Landscape Units (SELU) are produced in turn from LCFU and other geographical dimensions such as relief, belonging to a river basin, or proximity to the sea. LCFU are agglomerated with a methodology which maps dominant land-cover types. Large forests or agricultural areas will constitute a SELU in their own right while smaller units will be part of a larger zone characterised by its dominant land cover. The Dominant Land Cover Types are then classified according to river basins and relief classes (e.g. coastal, lowland, highland, mountain). The final intersection gives the map of terrestrial SELU.

Rivers are processed separately for accounting purposes. Rivers are land-cover units of a particular type where the dynamics of the water flow is the essence. In the case of rivers, the SELU will be the river system of the sub-basin. SELUs will be decomposed into drains (main drains, secondary drains) and segments (reaches) of homogenous water discharge.

In the case of seas, a distinction is made between the coastal zone, which is described as a 'seascape' that includes seabed features (as far as possible in conjunction with the coastal landscape). 'Open sea' is mapped according to various zonings, starting with fishery management areas (4).

The concept of **ecosystem services** can be found in the literature as far back as 1972 (Long, 1972). It was revitalised in the 1990s (Costanza, DeGroot, Daily...) and broadly used in the Millennium Ecosystem Assessment of 2006. Ecosystem services are defined as 'the contributions that nature makes to human well-being' (MA, 2006). In the case of the provisioning services, biomass products are usually measured in tonnes of carbon and water in cubic metres. The functional services, which are very heterogeneous, are measured as attributes of spatial units and weighted when relevant by population data. Here again, the concept of accessibility helps to switch focus from ecosystem functions to human well-being. The following definition is therefore proposed for accounting: ecosystem services are the outcome of ecosystem functions which are accessible to people.

2.1.2 Classifications

Draft classifications have been established, discussed in international meetings and will be submitted by the UNCEEA to global consultation. SECA implementation is based at this stage on the classifications in use in Europe.

The classification of Land Cover Functional **Units (LCFU)** is being prepared jointly by FAO and the EEA. It is based on the multi-purpose classification of homogenous land-cover types adopted for SEEA volume 1 and consists of the development of an application fit for classifying real landscape units which are more or less heterogeneous or mixed at the scale of SECA implementation. The documentation of LCFU will use FAO's Land Cover Classification System version 3 which has been acknowledged as a standard in the SEEA revision. It will define acceptable thresholds of heterogeneity and, in the case of mixed landscapes, the rules by which land-cover types combine to produce a new LCFU class (e.g. agriculture-nature mosaics or natural mosaics of shrubs and grassland). The forthcoming LCFU standard will easily bridge with the European Corine Land Cover classification, which is currently used for SECA. The common reference to LCCS3 will also enable the development of variants of the core LCFU at different scales and/or in particular geographical

Table 2.1 Land-cover types (first level) in SEEA volume 1

01	Artificial surfaces (including urban and associated areas)
02	Herbaceous crops
03	Woody crops
04	Multiple or layered crops
05	Grassland
06	Tree covered area
07	Mangroves
08	Shrub covered area
09	Aquatic or regularly flooded shrubs and/or herbaceous vegetation
10	Sparsely vegetated natural areas
11	Terrestrial barren land
12	Permanent snow and glaciers
13	Inland water bodies
14	Coastal water bodies and inter-tidal areas

Source: FAO, 2011.

⁽⁴⁾ More on statistical units for ecosystem accounting can be found in: Note on Statistical Units for Ecosystems, Alessandra Alfieri, Daniel Clarke and Ivo Havinga, United Nations Statistics Division and Jean Louis Weber, European Environmental Agency, Expert Meeting on Ecosystem Accounting, 11–13 May 2011, European Environmental Agency, Copenhagen, Denmark http://unstats. un.org/unsd/envaccounting/seearev/meetingMay2011/lod.htm.

Table 2.2 Aggregated Corine Land Cover used in Europe for LEAC and SECA

1	Artificial surfaces
2A	Arable land & permanent crops
2B	Pastures & mosaic farmland
3A	Forests and transitional woodland
3B	Natural grassland, heathland, sclerophylous (a) vegetation
3C	Open space with little or no vegetation
4	Wetlands
5	Water bodies

Note: (a) Vegetation that flourishes in hot dry areas.

Source: EEA, 2006.

Table 2.3 Provisional land-cover flow classification used in Europe for SECA

	nd development processes, urban sprawl, expansion nsive land use
lf11	Artificial development over agriculture
lf12	Artificial development over forests
lf13	Artificial development of other natural land cover
lf14	Conversion from small field agriculture and pasture to broad pattern cropland
lf15	Conversion from forest to agriculture
lf16	Conversion from marginal land to agriculture
lf17	Water body creation and management
lf2 Lar	nd restoration processes
lf21	Conversion from crops to set aside, fallow land and pasture
lf22	Withdrawal of farming
lf23	Forest creation, afforestation of agriculture land
If3 Rot	ations, natural processes and steady state
lf31	Internal conversion of artificial surfaces
lf32	Internal conversion between agriculture crop types
lf33	Recent tree clearing and forest transition
lf34	Forest conversions and recruitment
lf35	Changes of land-cover due to natural and multiple causes
lf4 No	observed land-cover change

contexts requiring more detail, while keeping the overall classification consistency.

- The classification of ecotones (the zones between major ecological communities) is derived from LCFU.
- The draft Classification of Land Cover Flows (LF) used in SECA is derived directly from the classification defined and used in the Land and Ecosystem Accounts (LEAC) report of 2006, the data of which were updated in 2011.
- A provisional classification of Socio-Ecological Landscape Units (SELU) has been established for test and discussion. The version below (Table 2.4) builds upon Corine land cover and LEAC methodologies for the definition of dominant land-cover types and relief classes. The classification of rivers and rivers basins is taken from the EEA's ECRINS database.

Table 2.4 Provisional classification of Socio-Ecological Landscape Units (SELU)

1. Mountai	in ecosystem landscapes
1.1	Urban and associated developed areas
1.2	Broad pattern agriculture
1.3	Agriculture associations and mosaics
1.4	Pastures and natural grassland
1.5	Forest tree cover
1.6	Other dominant natural land cover
1.7	Composite land cover (no dominant land cover)
2. Highlan	d ecosystem landscapes
2.1	Urban and associated developed areas
2.2	Broad pattern agriculture
2.3	Agriculture associations and mosaics
2.4	Pastures and natural grassland
2.5	Forest tree cover
2.6	Other dominant natural land cover
2.7	Composite land cover (no dominant land cover)
3. Lowland	d ecosystems (inland) landscapes
3. Lowland 3.1	d ecosystems (inland) landscapes Urban and associated developed areas
3.1	Urban and associated developed areas
3.1 3.2	Urban and associated developed areas Broad pattern agriculture
3.1 3.2 3.3	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics
3.1 3.2 3.3 3.4	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland
3.1 3.2 3.3 3.4 3.5	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover
3.1 3.2 3.3 3.4 3.5 3.6 3.7	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover Other dominant natural land cover
3.1 3.2 3.3 3.4 3.5 3.6 3.7	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover Other dominant natural land cover Composite land cover (no dominant land cover)
3.1 3.2 3.3 3.4 3.5 3.6 3.7 4. Coastal	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover Other dominant natural land cover Composite land cover (no dominant land cover) landscapes
3.1 3.2 3.3 3.4 3.5 3.6 3.7 4. Coastal 4.1	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover Other dominant natural land cover Composite land cover (no dominant land cover) landscapes Urban and associated developed areas
3.1 3.2 3.3 3.4 3.5 3.6 3.7 4. Coastal 4.1 4.2	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover Other dominant natural land cover Composite land cover (no dominant land cover) landscapes Urban and associated developed areas Broad pattern agriculture
3.1 3.2 3.3 3.4 3.5 3.6 3.7 4. Coastal 4.1 4.2 4.3	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover Other dominant natural land cover Composite land cover (no dominant land cover) landscapes Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics
3.1 3.2 3.3 3.4 3.5 3.6 3.7 4. Coastal 4.1 4.2 4.3 4.4	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover Other dominant natural land cover Composite land cover (no dominant land cover) landscapes Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland
3.1 3.2 3.3 3.4 3.5 3.6 3.7 4. Coastal 4.1 4.2 4.3 4.4	Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover Other dominant natural land cover Composite land cover (no dominant land cover) landscapes Urban and associated developed areas Broad pattern agriculture Agriculture associations and mosaics Pastures and natural grassland Forest tree cover

Provisional Common International Classification of Ecosystem Services (CICES)

In December 2008, EEA, together with UNEP and the German Federal Ministry of Environment, convened an international expert meeting on the project of a Common International Classification of Ecosystem Services (CICES). The need for such a standard results from the multiple global initiatives related to assessment and accounting of ecosystem services such as IPBES (Intergovernmental Platform on Biodiversity and Ecosystem Services), TEEB, MA follow-up, the European ecosystem assessment (Eureca!2012), many national assessments, Green Economics, PES and IPES (Payments and International Payments for Ecosystem Services), SEBI2010,

the SEEA2003 revision and the European Strategy on Environmental Accounting. CICES is expected foster synergies and bring together the diverse approaches taken to quantify and value ecosystem services.

Discussions took place at two international workshops on CICES hosted by the EEA in Copenhagen in December 2008 and 2009 and in an e-forum from November 2009 to January 2010, designed to enable a wider international audience to comment on the issues relating to the CICES concept. CICES was presented for information to the UNCEEA meeting of June 2010. The consultation is continuing.

CICES has been cross-referenced with CPC, the UN Common Products Classification.

Table 2.5 Provisional common international classification of ecosystem services

Theme	Class	Group
Provisioning	Nutrition	Terrestrial plant and animal foodstuffs
		Freshwater plant and animal foodstuffs
		Marine plant and animal foodstuffs
		Potable water
	Materials	Biotic materials
		Abiotic materials
	Energy	Renewable biofuels
		Renewable abiotic energy sources
Regulation and maintenance	Regulation of wastes	Bioremediation
		Dilution and sequestration
	Flow regulation	Air flow regulation
		Water flow regulation
		Mass flow regulation
	Regulation of physical environment	Atmospheric regulation
		Water quality regulation
		Pedogenesis and soil quality regulation
	Regulation of biotic environment	Lifecycle maintenance and habitat protection
		Pest and disease control
		Gene pool protection
Cultural	Symbolic	Aesthetic, Heritage
		Religious and spiritual
	Intellectual and Experiential	Recreation and community activities
		Information & knowledge


2.1.3 *Grids*

Although not statistical units in their own right, grids must be mentioned as important features of simplified ecosystem capital accounts. In Europe,

the INSPIRE Regulation defines a standard grid which is highly important when combining data from very diverse sources is needed — a constant in ecosystem accounting — or when analysis requires different scales or geographical breakdowns. In

Consumption at Full Price (including NDCEC)

Figure 2.1 Simplified ecosystem capital accounting structure (SECA)



[N] Ecosystem Monetary Balance Sheet: Stocks and Change of Ecosystem Financial Assets and Liabilities, in Euro

addition, grid data (raster) are faster to compute with GIS than vector data. The most widely-used grid for accounting purposes is $1 \text{ km} \times 1 \text{ km}$. The $0.1 \text{ km} \times 0.1 \text{ km}$ (1 ha) grid can also be used, but much less data is available at this scale.

2.2 The accounting structure of simplified ecosystem capital accounts

Simplified ecosystem capital accounts include tables in both physical and monetary units. Some of these tables are directly connected to SEEA volume 1 tables where breakdowns are mostly presented by economic sector and are, in that way, indirectly bridged to the SNA itself (in particular regarding supply and use and input-output tables). Other tables link back directly to the SNA.

The following sections comment on the various tables one by one. The tables are presented separately in a spreadsheet with mock-up numbers aiming to facilitate ease of understanding (see Annex 2). The accounting structure is summarised in Figure 2.1.

2.3 The physical accounting tables

2.3.1 The basic balances

Table [A] Land-cover stocks and flows basic account

This account measures, in km², the land-cover stocks and changes in the ecosystem statistical units used for accounting.

Land-cover stocks: artificial surfaces, large to medium farm arable land and permanent crops, pastures, mosaic farmland (small farms, mixed land cover), forest cover, natural grassland, scrubland, natural mosaics, open space with little or no vegetation, wetlands and water bodies.

Elementary one-to-one land-cover changes are grouped into land-cover flows: land development processes, urban sprawl, land-use intensification, land restoration processes, rotations, natural processes and steady state.

Land-cover flows are **indicators of land-cover consumption** (regarding the opening year) **and new formation** (in the closing year).

Produced from satellite images, land-cover accounts can be reported at various scales, regarding various types of natural or administrative zonings and by regular grids (1 x 1 km or 0.1 km x 0.1 km). Their organisation plays a central role in organising the whole system of inland ecosystem accounts.

Table [B] Ecosystem capital carbon/biomass account

The ecosystem capital carbon/biomass account measures the Net Ecosystem Accessible Carbon Surplus (NEACS) in soil, vegetation and fisheries and its use.

The account records, in tonnes of carbon, the stocks available in soil, below-ground and above-ground vegetation and in water (fish). It records the flows of Net Primary Production (NPP) by natural and cultivated vegetation, and its use by crops and timber harvests. In addition to inland ecosystems, the accounts covers sea (fisheries and sea regulating capacity) and the atmosphere's climate regulation capacity which is a measure of the amount of fossil carbon accessible without increasing mean global temperature beyond the stated target of a maximum of 2 degrees Celsius.

The characteristic indicators of ecosystem capital carbon/biomass accounts are:

- NPP and its removal by agriculture, forestry and fisheries, which indicates the availability of these provisioning ecosystem services;
- the Net Ecosystem Carbon Balance (NECB)
 which indicates the sustainability of carbon/
 biomass use; in principle, NECB should be
 always greater than or equal to zero;
- Net Ecosystem Accessible Carbon Surplus
 (NEACS) which measures the share of available
 ecosystem production of carbon which meets
 the sustainability constraints of maintaining
 stocks in soils and vegetation (mostly in trees)
 and fisheries; in addition to inland and sea
 ecosystems, NEACS includes the fossil carbon
 accessible under constraint of maintenance of
 the atmosphere's climate regulation functions.
- The Ecosystem Accessible Carbon Surplus index summarises the sustainability of total carbon use (removal of biological carbon plus use of fossil carbon) compared to the accessible resource (NEACS). The ratio NEACS/Use should be always greater than zero.

Because of the primary character of biomass ecosystem production and the comprehensive coverage of carbon/biomass accounts, they play a central role in ecosystem capital accounts. Carbon/biomass is the primary service expected from the ecosystem, under the constraint of sustainable supply of water (for human use and the ecosystem itself) as well as sustainability of all the regulating (water regulation, assimilation of residuals, habitat regulation, pollination) and socio-cultural services which are produced.

Carbon/biomass use and ownership can also be detailed by economic sectors, which is done in another part of the SEEA, so-called the 'supply and use tables' and 'asset accounts'. Carbon/biomass can in that way be brought together with the SNA tables in money terms ('supply and use' and 'input-output') for hybrid analysis and modelling.

The use of biological and fossil carbon can be or is recorded in national accounts by economic sectors and commodity content (embedded carbon). Carbon/biomass accounts are therefore an essential element to broaden the scope of **resource efficiency indicators** towards integrating impacts on the ecosystem (the 'second decoupling' paradigm).

Table [C] Ecosystem Capital Water Account

The account measures the Total Ecosystem Accessible Fresh Water (TEAW) and the Net Ecosystem Accessible Fresh Water Surplus (NEAWS) adjusted for water stress during the vegetation growing season.

Accounts in m³ are established for water stocks in terrestrial ecosystems (soil and vegetation) and water bodies (aquifers, lakes and dams, rivers). They include a distinction between total and accessible stocks, the difference being due to physical or economic constraints of abstraction, pollution or time mismatch between availability and requirements for natural or human uses.

The water flow accounts are tracked from precipitation infiltration and runoff down to the final outflow. Total available *effective* rainfall (in hydrological terms), which is available to feed the water bodies, is precipitation minus Evapo-Transpiration (ETa). ETa is subdivided into 'spontaneous' and 'induced by irrigation and other uses'. 'Spontaneous' ETa is further subdivided into 'induced by rain-fed cultivated vegetation' and 'induced by non-cultivated vegetation'. Total available effective rainfall is further analysed to

take account of inaccessible water due to events like floods, wastewater disposal and dilution, additional ETa induced by irrigation, and evaporation induced by power plants cooling towers or reservoirs. Total ecosystem accessible fresh water can then be computed after appropriate adjustments to take account of water transfers between ecosystems and within or between river basins. A final adjustment is then made to reflect the timeliness of the water resource regarding vegetation requirements.

The characteristic indicators of water accounts are:

- Total available effective rainfall, calculated from a hydrological perspective (water available for runoff), before evapo-transpiration induced by irrigation and evaporation induced by other uses;
- Withdrawals of water (by ecosystems, catchments and economic sectors);
- Returns of wastewater, an additional although degraded resource and a cost regarding maintenance of ecosystem water quality. Taking into account the acceptable dilution of pollutants (maximum concentration, BOD), untreated wastewater returning to water bodies can reduce the accessibility of fresh water by several times the amount of wastewater discharged;
- Returns of water to soil due to losses in transport and irrigation;
- Total Ecosystem Accessible Water: TEAW is the accounting balancing item of stocks and flows;
- Water Stress Coefficient is an additional adjustment needed to reflect the timeliness over the year of water accessibility considering vegetation requirements, in particular during the growing period;
- Net Ecosystem Accessible Fresh Water Surplus.

The final aggregate is called **Net Ecosystem Accessible Fresh Water Surplus (NEAWS)**. It can be compared to the withdrawals of freshwater to measure the intensity of use of the water resource. The ratio Withdrawals/NEAWS should always be < 1. A lower target value is likely to be needed in order to take stock of the variability of the water resource and the economic and social acceptability of risks of periodic deficits and thus the sustainability of the withdrawals.

Table [D] Landscape green infrastructure accounts

These accounts measure the capacity or potential of ecosystems to deliver ecosystem services in a sustainable way. Typical indicators are Landscape Ecosystem Potential (LEP), Green Accessible Landscape Infrastructure (GALI) and Rivers Ecosystem Potential (REP). They reflect the fact that sustainable provision of carbon/biomass and water has to be compatible with the good functioning of ecological infrastructure, landscapes and rivers and that access to the many ecosystem regulating and socio-cultural services is better measured in the first instance by the abundance and health of the ecosystems which deliver them to people. Ecosystem health is assessed following the principles developed by David J. Rapport in his description of the 'ecosystem distress syndrome' (Rapport 2007).

In contrast to provisioning services, regulating and cultural services cannot be harvested or abstracted and consumed. Their value in the economy is seen as an attribute of land which is not measured directly but bound up with the values of real estate and/or related goods and services. Disentangling such ecosystem services from market values has been done in many case studies for selected services. However, there is as yet no evidence that such case studies can be up-scaled to the macro level of national accounts. Service-specific accounts have been generated successfully on a one by one basis but there is no evidence that double counting when aggregating several of these services can be avoided or that the full range of ecosystem services can be covered. Simplified ecosystem capital accounts therefore start by measuring, in a holistic way, the capacity of the capital to continue to deliver any service over time. This approach does not preclude the development of local and/or service-specific accounts but offers instead a starting point and background information for such exercises.

The characteristic indicators of landscape green infrastructure accounts are:

- The Green Background Landscape (GBL) index, which weights hectares of land cover according to their 'greenness'. Because the greenness of an ecosystem service is important not only in each place but also in its neighbourhood, the calculation is based on fuzzy logic (smoothed values);
- The Mesh Effective Size (MEFF) index, which measures the partitioning effects of landscape fragmentation by urban areas and transport

- infrastructures. Small meshes limit good ecological functioning;
- The Stated Social Nature Value (SSNV) index, which reflects the importance of biodiversity for society as expressed in terms of landscape protection;
- Landscape Ecosystem Potential (LEP) a multi-criterion index which combines GBL, MEFF and SSNV;
- The Green Ecotones Index, based not on the surface area of land-cover units but on the length of their borders which concentrate higher animal and plant biodiversity. The ecotone classes are weighted according to their potential for hosting biodiversity;
- The Green Accessible Landscape Infrastructure index (GALI) is a composite index which combines GBL and the green ecotones indexes;
- The River Infrastructure Potential, which measures the capacity to deliver water and related services of large, medium and small rivers and brooks and streams; it is measured in a common unit named 'standard-river km' (1 srkm = 1 km x 1 m³/second);
- The River Integrity Composite index, which combines indexes of water quality, river fragmentation and river green ecotones.
- Rivers Ecosystem Potential (REP), a composite index combining river infrastructure potential and integrity. REP connects landscapes to water accounts.

All indexes are produced from spatial analysis and assimilated into a regular grid (typically 1 km² cells at the macro level, 1 hectare at the local level) which facilitates the detection of interactions, potentials and degradation. In a second step, elementary and/or composite indexes can be aggregated by regions, catchments etc.

Table [E] Ecosystem Capital Biodiversity Account

This account brings together biodiversity variables measured from the standpoints of landscape and species/biotopes. The first sub-account of **Biodiversity Infrastructure Integrity (BII)** is a continuation of Table [D] from which it is computed. It is supplemented by a second sub-account based

on species and biotope monitoring which is used for producing a diagnosis of ecosystem health or distress (Rapport, 1999). Several methods can be accepted for that, the aim being to assess ecosystem health regarding biodiversity, not biodiversity per se. Finally, the Ecosystem Biodiversity Rating (EBR) combines landscape and species/biotope indexes.

2.3.2 Synthesis tables in physical units: Total Ecosystem Capital Potential and Physical Balance Sheet (Assets and Liabilities)

The ultimate purpose of ecosystem capital accounts is to assess whether economic use of ecosystems results in an increase and/or improvement, a steady state, or a degradation of the natural assets which together are used as economic resources, consumed by the economy as free externalities, and/or directly supply a range of free services to individuals or humankind as a whole. As the measurement of ecosystem capital on the basis of private benefits is necessarily incomplete, it can be misleading. Ecosystems are multifunctional and the core of the issue is that using one particular function or service most often results in the degradation and even elimination of one or all the others. The other possible approach is to look at the capital as a bio-physical system and assess its ability to continue to deliver its services.

As noted above, the SECA model considers three groups of services: biomass/carbon production, freshwater production and functional services. It measures, for each of them, the amount that is accessible regarding the (maximum) accessible surplus that can be used without impairing the reproduction of nature itself and the need to ensure that the use of one of these services does not degrade access to the others. For each group, it is an issue of maximisation of the resource in the presence of internal and external constraints. As the three assets and their services are not measured in additive units, one has to be chosen as the primary one. The proposal is that this should be biomass/carbon, the primary component of life, encompassing food, fibre and energy. The narrative of SECA is then as follows: ecosystem capital is measured by the stocks of accessible biomass/carbon adjusted in the light of constraints of freshwater accessibility, maintenance of landscape and river potentials and biodiversity conservation.

Table [F1] Total Ecosystem Capital Potential Account

This table presents the calculation of Net Total Ecosystem Capital Potential (NTECP) and Net Change and Territorial Ecosystem Capital Degradation (TECD). The starting point is given by table [B]. The balancing item 'Net Ecosystem Accessible Carbon Surplus' (NEACS) is taken as a surrogate measure of the gross ecosystem capital potential of inland, sea and atmosphere ecosystems. In simplified accounts it covers the accessible carbon of terrestrial ecosystems, sea (fisheries) and the atmosphere's capacity to assimilate carbon. Estimation of similar assimilation capacity should be done for the sea. River system potential also needs to be measured in a consistent way and added to the gross total potential. A possible solution is to assess the exergy (accessible energy) potential of rivers following the approach to water accounting in Spain developed by Naredo, Valero et al. (Valero, 2006). An important point is that this work starts from accounts of river stocks which are identical to the River Infrastructure Potential of Table [C] in terms of 'standard-river km' (1 srkm = $1 \text{ km x } 1 \text{ m}^3$ / second). The solution would be to convert 'srkm' measurements into exergy potential and then into carbon unit-equivalents.

At this stage, 'gross' means prior to integration into the calculation of the other factors that limit the accessibility of the carbon resource: access to other services and maintenance of ecosystem functions. This integration will be done by weighting the gross potential with indexes extracted from tables [C], [D] and [E]: Ecosystem Accessible Water Surplus (EAWS), Landscape Ecosystem Potential (LEP), Rivers Ecosystem Potential (REP) and Ecosystem Biodiversity Rating (EBR). Several options are available for combining the various indexes into a 'limiting factors index' to weight the gross potential, including average values or more elaborate methods e.g. Bayesian belief network decision trees. The final decision will have to take into consideration that it is more important to quantify change than to quantify stocks; sensitivity analysis will be necessary to establish the final methodology.

By weighting the initial carbon balance with the limiting factors index, we create a measurement unit of general application which we will call Ecosystem Potential Unit Equivalents (EPUE).

Within an economic territory (using the SNA definition), Net Total Ecosystem Capital Potential (NTECP) can increase or decrease. An increase reflects ecosystem improvement due to restoration programmes or spontaneous natural processes. A decrease can be the effect of natural disturbances or ecosystem degradation. Ecosystem Capital Degradation (ECD) is thus defined in a strict way as the consequence (effect, impact) of human activities.

Table [F2] Account of Territorial Ecosystem Capital Degradation (TECD) by Stress Factors

Table [F2] analyses TECD by stress factors.

Stress factors are: effect of land-cover change, restructuring/de-structuring of landscapes and rivers, over-exploitation of biological resources, waste disposal and pollution (including GHGs). The link to sectoral accounts is through the rows of supply and use tables detailing the generation of pollutants and emissions of residuals and the more elaborated hybrid input-output tables (5) (combining physical and monetary data). Other links are with land-use accounts which bridge to agriculture and forestry statistics of crop yields and farming and management practices, and with fisheries accounts and statistics.

Table [F2] is used later when calculating ecosystem capital depreciation (see tables [I] and [J]).

Table [H] Physical Balance Sheet: Assets and Liabilities

The Ecosystem Capital Physical Balance Sheet brings together the physical ecosystem assets (from Table [F1]) and the physical debts or liabilities that the economy contracts to future generation when degrading nature. This concept of physical liability does not exist in the SNA where both financial and non-financial assets are balanced by debts which are all recorded in the financial tables. This practice conforms to the analysis of the economic system. In ecosystem accounts, as long as some costs are not paid by the economy, it is necessary first to record the physical degradation as a liability. Then, Table [H] is supplemented with a second balance sheet in money terms. This prevents changes in natural asset value being seen as resulting in an improvement in the situation when physical degradation is not remediated or compensated.

The physical balance sheet is established in Ecosystem Potential Unit Equivalents (EPUE) — see Section 1.2.

It presents:

- physical ecosystem assets accounts where the opening balance is the initial Net Total Ecosystem Capital Potential; Change in Ecosystem Potential Due to Economic Activities, Other Change in Potential of Ecosystem Capital make Net Change in Physical Ecosystem Assets leading to the closing balance (the resulting NTECP);
- physical liabilities accounts include, as the opening balance, NTECP plus international national and private Ecosystem Restoration Targets (recovery from historical damage, compliance with conventions/regulations) endorsed by society. Changes are due to Territorial Ecosystem Capital Degradation (TECD), Ecosystem Capital Degradation in 'consumed imports', and, in the opposite direction, reduction of physical liabilities by ecosystem restoration programmes and spontaneous natural improvement of ecosystems, reduction of physical liabilities by acquisition of EPUE (mitigation/compensation), and reduction in physical liabilities by swaps and debt consolidation. Opening balance sheet plus Net change in physical liabilities makes the Closing balance sheet item for physical liabilities.

Table [G] Demand for and Accessibility to Ecosystem Services

Table [G] details the demand for and accessibility to Carbon/Biomass, Fresh Water, and Green Infrastructure Neighbourhood Ecosystem Services (GINES). The tables are presented by ecosystem type and include population data in order to calculate accessible ecosystem services per capita. Accessible Carbon/Biomass and Accessible Fresh Water were presented in Tables [B] and [C] respectively. When referring to local or gridded population data, the indicators represent accessibility in the place (ecosystem or grid-cell) or in the neighbourhood when using fuzzy datasets (e.g. data smoothed over a radius of 5 km) (6).

⁽⁵⁾ Known in Europe under the acronym of NAMEA.

⁽⁶⁾ See EEA, 2006.

Green Infrastructure Neighbourhood Ecosystem Services (GINES) is derived from Tables [D] and [E]. It is an over-arching indicator which assumes a direct relationship between ecosystem health and the availability of regulating and cultural ecosystem services. Its calculation takes into account the fact that landscape artificial intensity reduces the supply of green infrastructure services and, at the same time, increases the number of potential beneficiaries (because of neighbourhood or easier access by transport infrastructures). Not surprisingly, medium-size human settlements will be the best performers regarding accessible GINES.

GINES are not only interesting per se. As long as they can be computed by Socio-Ecological Landscape Units and/or grid-cells and reported by river basins and administrative units, GINES can be balanced with other variables in trade-off analysis. The first of these are Accessible Carbon/biomass and Accessible Fresh Water as well as Net Total Ecosystem Capital Potential and Territorial Ecosystem Capital Degradation.

2.4 The monetary accounting tables

Monetary accounts for ecosystem capital and services will be developed on top of physical accounts. As noted by the UNCEEA in June 2011, not all possible valuation methods are relevant to national accounting, only those that are compatible with the SNA rules.

Compatibility with the SNA excludes some methods frequently used in cost-benefit analysis (typically 'contingent valuation') because of different definition of value itself (based on observable transaction prices in the SNA, on willingness to pay in CBA) and of up-scaling and aggregation issues (Weber, 2011a). It does not, however, exclude the estimation of important economic variables in the absence of directly observable transactions. This is the case for 'government services' which are valued by the total of production costs (but exclude any operating surplus), the production of food products for own use (value using the basic price in farms, not at purchaser price of similar products), etc.

One particularly interesting variable which has to be estimated is Consumption of Fixed Capital (CFC), the equivalent in SNA to capital depreciation in business financial accounting. CFC is the accounting item which makes the difference between the Product and Income concepts. According to the International Accounting Board standard, capital depreciation *must* be subtracted when calculating a

company's profit. The point is that consumption of ecosystem capital is not subtracted in the same way, because it is considered as an externality, a cost to be borne by society, not by those responsible for the ecosystem degradation. It is an unpaid cost which biases the estimation of growth and progress given by the conventional national accounts aggregates: Net Domestic Product or National Income — which are overestimated, and Final Consumption (and Imports) — which are underestimated, leading to the well-known distortions of consumption patterns and international trade (Chichilnisky, 1994).

Paying for ecosystem capital depreciation is however an idea at work in several areas:

- A prime example is the Clean Development
 Mechanism of the Convention on Climate
 Change which is based on accounting for carbon
 and CO₂. The target of 'maximum temperature
 increase of 2 degrees' refers to degradation
 of the atmosphere ecosystem. The cost of
 keeping below this target is 'ecosystem capital
 consumption'.
- Reducing Emissions from Deforestation and forest Degradation (REDD) was originally framed as payments for a particular ecosystem service (carbon sequestration by forests) and has moved to REDD+, which considers ecosystem capital degradation more holistically (to avoid biological and other leakages).
- Another example is given by the European Environmental Liability Directive of 2004 and the shift in the Polluter Pays Principle from pressures towards ecosystem impacts. The remediation costs of these impacts are ecosystem capital consumption.
- A similar approach has been taken in the European Water Framework Directive with the over-arching targets of 'good environmental quality of the river basins' (to be quantified and for which costs of remediation measures have to be estimated by Member States) and 'full recovery of costs' in water pricing.
- Wetlands Mitigation Banking in USA is another example where 'accounts' are established for 'ecosystem service areas' defined as 'the designated geographic area in which a bank can reasonably be expected to provide appropriate compensation for unavoidable impacts to wetlands.' The 'determination of credits' in a 'wetland bank' is based on accounts of physical characteristics and capacity to deliver services

which 'include acreage, category type, and/or function'. The need for credits for mitigating damage is assessed by a certification process based on a symmetric accounting of the amount to be replaced regarding expected damages. This is the basis of a market mechanism where credit values are established.

 Other examples can be found in timber certification mechanisms and last but not least in the expanding 'fair trade' market voluntarily supported by more and more consumers. We should pay the real price of what we consume.

Till now, integrating unpaid costs into retrospective national accounts has been rejected by national accountants on the grounds that it would implicitly modify the prices-consumption structure which has led to a particular GDP amount. We cannot re-write the past (which is observed by statisticians) and such adjustments should be envisaged only for modelling the future (Bosch, Brouwer, Radermacher et al.,1997; or Vanoli, 2005).

While it is certainly not possible to modify the past as recorded in national accounts, there is a solution for taking stock of unpaid consumption which is to record it as a debt. For that purpose, monetary accounts for ecosystem capital include a detailed financial balance sheet.

[I] Estimation of unit costs of ecosystem capital restoration by stress factors

The estimation follows the structure of Table [F2] of physical degradation by stress factors. In Table [I], unitary costs per EPUE are derived from analysis of real expenditures or costs of restoration programmes. Such work relies on the expertise of environmental agencies, water agencies, agronomists, foresters, etc, doing such calculations in their daily work. Estimates of unitary costs have to be carried out by ecosystem types/issues/regions.

[J] Territorial Ecosystem Capital Depreciation

Table [J] presents the estimation of **Territorial Consumption of Ecosystem Capital (TCEC)** in money. TCEC is calculated as degradation in EPUE multiplied by unitary remediation costs by ecosystem types/issues/regions.

[K] Accounts of Ecosystem Capital Degradation & Depreciation Embedded in Imports & Exports

The accounts of the SNA are compiled for resident institutional units grouped into institutional sectors and subsectors. Together these describe the Domestic rather than National (used for qualifying Income) or Territorial economy (which applies to the physical world). An institutional unit is resident in a country when it has a centre of economic interest in the territory of that country. Resident units may have temporary activities in the rest of the world, and may import and export commodities. In terms of ecosystem capital degradation, this means that the territorial approach should be broadened to take account of the effects of the domestic economy on the rest of the world, in particular from the use of natural resources and the degradation that this may generate in the producing country. This can be due to uneven level of environmental protection or uneven robustness of property systems (Chichilnisky, 1994) which results in flows of ecosystem capital consumption embedded in international trade (Koellner, 2011). These embedded flows must be measured and valued in an appropriate way, which is the purpose of Table [K]. Virtual or embedded land in 'consumed imports' (Van der Sleen, in Koellner, 2011) is recorded first as data infrastructure, then as ecosystem capital degradation embedded in imports of agriculture, forest, and fishery products and carbon embedded in the production of all imported products. Ecosystem capital degradation embedded in imports (in EPUE) is then converted into 'unpaid ecosystem depreciation' in 'consumed imports' in money terms. At this stage, remediation costs should be estimated on the basis of prices in the importing country.

[L] Sustainable Macro-economic Benefits from Ecosystem Services

The macro approach chosen for simplified ecosystem capital accounts in Europe results in a service-by-service assessment of selected benefits. Ad hoc calculations can be well supported by the physical accounts presented above. Some of these can be done for specific services and integrated into the SECA framework. We propose to start with provisioning services, following recent experiences in Zanzibar (Lange & Jiddawi, 2009) and for the Mediterranean Sea (Plan Bleu, 2010). The methodology currently tested at the EEA (7)

⁽⁷⁾ Acosta, J., Pedersen, O. G. and Weber, J.-L., 'Ecologically Sustainable Total Induced Value Added', draft, EEA and European Topic Centre on Sustainable Consumption and Production (ETC-SCP), Copenhagen, 2011.

is called 'hypothetical extraction' of value-added from input-output tables (I-OT). The I-OT are those compiled for Europe by Eurostat for 1995 to 2005. The first result expected is the Total Induced Value Added of key sectors, starting with agriculture and the food industry, followed by forestry, fisheries and water management. In a second step, TIVA is planned to be adjusted with an index of ecosystem capital degradation for the respective sectors, leading to Ecologically Sustainable Total Induced Value Added (ES-TIVA). Note that the ES-TIVA for the various sectors cannot be added together without double counting.

[M] Economic aggregates and additional adjustments for Consumption of Ecosystem Capital (CEC)

Table [M] takes over the conventional SNA aggregates and the Consumption of Fixed Capital. It appends the summary of ecosystem capital calculation: Territorial Consumption of Ecosystem Capital plus Ecosystem Capital Depreciation virtually embedded in imports equals Gross Domestic Consumption of Ecosystem Capital (GDCEC). GDCEC minus Ecosystem Capital Depreciation virtually embedded in exports equals Net Domestic Consumption of Ecosystem Capital.

Finally, Table [M] proposes additional macroeconomic aggregates adjusted for ecosystem depreciation. They are GDCEC Adjusted Net Domestic Product or National Income, Final Consumption at Full Price (which includes NDCEC), Imports at Full Price and Export at Full Price.

[N] Monetary Balance Sheet: Assets and Liabilities

Table [N] is the Monetary Balance Sheet of ecosystem Assets and Liabilities. To a large extent, Table [N] mirrors Table [H], the Physical Balance Sheet. The two separate balance sheets are necessary as long as SECA does not postulate that ecosystem capital degradation can automatically be compensated by an expenditure. Instead, SECA considers remediation costs as an estimate (in the way Fixed Capital Consumption is an estimate) of the amount which should be reinvested in the

next period to repair the observed degradation. If degradation persists despite restoration action, a new Consumption of Fixed capital will be calculated accordingly. The relative evolution of physical and monetary liabilities, in the territory and in the rest of the world, is likely to become an important assessment variable (8).

On the asset side, one must note that no calculation of total ecosystem capital potential monetary value is foreseen. The remediation costs represent the costs of restoring the ecosystem, which at some moment will be a natural process. Using remediation costs to calculate the value of the ecosystems themselves would be seriously misleading since it would imply that nature is produced by human activity — when it is at best a co-production. The only costs recorded are the market value of economic ecosystem assets and the financial assets which can be accumulated as a consequence of ecosystem improvement.

Conclusion

This paper is followed by annexes with details of the various tables described briefly above. One version is a mock-up table in which real, estimated and invented numbers have been used. This does not matter at this stage as long as it is appreciated that the numbers are there to enable understanding of the way the various accounting items relate to each other and are used to synthesis tables and aggregates. Simulated numbers are being progressively replaced by real ones, in the course of the fast-track implementation of the Simplified Ecosystem Capital Accounts in Europe. Not all details are currently there (it is still a simplified framework) and there is room for improvement. However, the present sketch gives a vision of possible accounts of the ecosystem in its co-evolution (9) with the economy.

The ecosystem capital accounting framework of Table 6 integrates physical and monetary tables. Physical tables integrate basic quantitative balances and qualitative indexes of ecosystem health and ecosystem services accessibility. Basic quantitative balances compiled by ecosystem type can be mirrored with the physical supply and use tables and economic assets accounts detailed by economic sectors in SEEA volume 1. In that way

⁽⁸⁾ It can be envisaged that ecological debts exported/imported with non-sustainable products are recorded by a special international institution. They would be recorded twice, in physical and in monetary units. Payments by the debtor would be made to this institution which would repay the creditor in proportion to the effective remediation of ecological damages.

⁽⁹⁾ To quote Norgaard, 1994.

they can be connected to the monetary Supply and Use Tables (S-UT) and natural assets accounts of SNA 2008 itself. Ecosystem capital accounts are mainly built up from geo-referenced data so they are genuinely top-down, connected to the local scales as much as to the macro level. A large part of the accounts can be reported by regions or river basins. Ecosystem capital accounts measure resource stocks and flows, factors limiting use, and the accessible resource surplus, and compare it with resource use computed from statistics. They measure ecosystem degradation, remediation costs and the accumulation of ecological debts which may result from cumulative degradation both in the country and abroad in trade-partner countries. All these elements are particularly important for monitoring progress towards green economy and assessing green growth. In particular they broaden the scope of the resource efficiency indicators based on material flow analysis: the flows can now cycle between ecosystems, and opportunities, quantitative and qualitative constraints and risks can be taken into account. Assessment of progress or degradation of well-being is no longer restricted to market variables.

The issue of how to handle subsoil assets, coal, oil and minerals remains. They are the 'dead

services' of ecosystems dead for hundreds of millions of years. They are not being renewed on any significant time scale that people can influence. Their depletion is primarily an economic issue. The way to integrate them into the picture in a way consistent with the physical/money approach of ecosystem degradation could be to follow El Serafy's 'user cost' method (El Serafy, 1992). This aims at measuring the share of the economic benefit from resource exploitation which should be reinvested in another asset in order to maintain the resource flow at a constant level. Another method for measuring economic asset depletion is proposed with variants by the SNA and SEEA volume 1; this should give similar results when prices are not too volatile. The two approaches however do not reflect the fact that, having alleviated the pressure on ecosystems at the beginning of the industrial revolution, subsoil energy and minerals have become by far the most important source of pollution and poisoning of ecosystems. A weak approach to sustainability of the ecosystem issue (i.e. maintain income, not the environment) is hence very debatable. It is an open question whether to expand the ecosystem capital accounting approach to fossil resources, on the basis of ecosystem assimilation capacity which finally determines their accessibility (10).

⁽ 10) It would follow the pioneer work of Naredo and Valero (Valero, 2006).

Bibliography

Blue Plan, 2000, Margat, J., Vallée, D., Water Resources and uses in the Mediterranean Countries: Figures and Facts, 2000.

Bosch, P., Brouwer, R., Radermacher, W. et al., 1997, *Methodological Problems in the Calculation of Environmentally Adjusted National Income Figures (GREENSTAMP Final Summary Report)*, Report for the European Commission Directorate General XII, Contract No. EV5V-CT94-0363, July 1997.

Chichilnisky, G., 1994, *North-South Trade and the Global Environment* (1994). American Economic Review, Vol. 84, No. 4, September 1994. Available at SSRN: http://ssrn.com/abstract=1374644.

Costanza, R., 1997, The Value of the World's Ecosystem Services and Natural Capital, Costanza, R., D'Arge, R., DeGroot R., Farber, S., Grasso, M., Hannon B., Limburg, K., Naeem, S., O'Neill, R., Paruelo, J., Raskin, R., Sutton, P. and van den Belt, Nature, M. 387, 253–260.

Daily, G. C., 1997, Introduction: What are Ecosystem Services? In: Daily, G.C. (Ed.) *Nature's Services: Societal Dependence on Natural Ecosystems*. Island Press, Washington, D.C., 1–10.

EEA, 2006, Land accounts for Europe 1990–2000, EEA Report No 11/2006, drafted by Haines-Young, R. and Weber, J.-L., http://reports.eea.europa.eu/eea_report_2006_11/en.

EEA, 2010a, Ecosystem Accounting for the Cost of Biodiversity Losses: The case of Coastal Mediterranean Wetlands, EEA Technical Report No 3/2010, drafted by Haines-Young R., Potschin M., Kumar P., Weber J.-L. et al., http://www.eea.europa.eu/publications/ecosystem-accounting-and-the-cost.

EEA, 2010b, Proposal for a Common International Classification of Ecosystem Goods and Services (CICES) for Integrated Environmental and Economic Accounting, Paper prepared for the EEA by Centre for Environmental Management, University of Nottingham, United Kingdom, Background document, ESA/STAT/AC.217, UNCEEA/5/7/Bk, Fifth Meeting of the UN Committee of Experts on Environmental-Economic Accounting, New York, 23–25 June 2010, http://unstats.un.org/unsd/envaccounting/ceea/meetings/UNCEEA-5-7-Bk1.pdf.

El Serafy, S., 1991, The Environment as Capital. 1991. Chapter 12, pp. 168-175 in Robert Costanza (editor), Ecological Economics — the Science and Management of Sustainability. Columbia University Press, New York.

El Serafy, S., 1992, Sustainability, Income Measurement and Growth. 1992. Chapter 5, pp. 63-79 in Robert Goodland, Herman E. Daly and Salah El Serafy (editors), *Population, Technology and Lifestyle: The Transition to Sustainability*. Island Press, Washington DC.

FAO and EEA, 2009, Land Cover and Land Use Classifications in the SEEA Revision, drafted by Xiaoning Gong (FAO) and Jean-Louis Weber (EEA), Fourth Meeting of the UNCEEA24 — 26 June 2009, UN Headquarters, New York, http://unstats.un.org/unsd/envaccounting/ceea/meetings/UNCEEA-4-11.pdf.

FAO, 2011, Towards a System of Environmental Economic Accounting for Agriculture (SEEA-AGRI), Paper prepared for FAO by Juan-Pablo Castañeda and Jaime Luis Carrera, Institute of Agriculture, Natural Resources and Environment, Rafael Landívar University, Guatemala, ESA/STAT/AC.238, UNCEEA/6/27, http://unstats.un.org/unsd/envaccounting/ceea/meetings/UNCEEA-6-27.pdf.

Friend, A. M., 2004, *Valuation of the Physical Stock-Flow Accounts through means of Ecopricing Sraffa Ecosystem Valuation Method SEVM*, The 8th Biennial Scientific Conference of the International Society for Ecological Economics, Montreal, 11–14 July 2004.

GAISP, 2006, 2010, Eight monographs (of which 3 related to forests) and a methodological note, Haripriya Gundimeda, Pavan Sukhdev, Pushpam Kumar, Rajiv Sinha and Sanjeev Sanyal, http://www.gistindia.org/publications.asp.

Gallopin, G.C., 1991, Human dimensions of global change: linking the global and the local processes. *International Social Science Journal* 43(4):707–718.

Glaser, M., Krause, G., Ratter B. and Welp, M., 2008, 'Human-Nature Interaction in the Anthropocene: Potential of Social-Ecological Systems Analysis', *GAIA*, Vol.17, No.1, pp.77–80.

Houdet, J., Trommetter, M., Weber, J., 2010, *Promoting business reporting standards for biodiversity and ecosystem services. The Biodiversity Accountability Framework.* Orée, 16 p., http://www.business-biodiversity.eu/default.asp?Menue=153&News=310.

IASB (International Accounting Standards Board), 2009, *International Financial Reporting Standards* 2009, ISBN: 978-1-905590-90-2, http://www.iasb.org/.

Koellner, 2011, *Ecosystem Services and Global Trade of Natural Resources*. Thomas Koellner Editor, Routledge 2011 – in particular Part 2: Ecosystem impacts of global flows of virtual land, water and sea in the physical economy, http://www.routledge.com/books/details/9780415485838/.

Lange G.-M. and Jiddawi N., 2009, 'Economic value of marine ecosystem services in Zanzibar: Implications for marine conservation and sustainable development', *Ocean & Coastal Management*, Volume 52, Issue 10, October 2009, pp. 521–532, http://www.sciencedirect.com/science/article/pii/S0964569109000994.

Commission interministérielle des comptes du patrimoine naturel, 1986, *Les comptes du patrimoine naturel*, Collections de l'INSEE, C137-138, Paris, 1986.

Long, Gilbert, 1972, *A propos du diagnostic écologique appliqué au milieu de vie de l'homme*, Options Méditerranéennes n°13, 1972, Centre International de Hautes Etudes Agronomiques Méditerranéennes (CIHEAM), http://ressources.ciheam.org/om/pdf./r13/CI010462.pdf.

MA [Millennium Ecosystem Assessment], 2006, *Ecosystems and Human Well-being: Synthesis*. Island Press, Washington, DC.

Martinez A., 2011, Assessment of Environmental Water Cost Through Physical Hydronomics, Amaya Martínez, Javier Uche, Antonio Valero, Carlos Rubio, Water Resources Management, DOI: 10.1007/s11269-011-9786-1, http://www.springerlink.com/content/m7rv6t144g838327/.

Naredo J. M., 2007, Costes y cuentas del agua. Propuestas desde el enfoque ecointegrador, Seminario Costes y Cuentas del agua en Cataluña en relación con la Directiva Marco del Agua, Agencia Catalana del Agua, 18–19 June 2007, http://www.fnca.eu/fnca/docu/docu/229.pdf.

Norgaard, R. B., 1994, *Development Betrayed: The End of Progress and a Coevolutionary Revisioning of the Future.* Routledge. London.

O'Connor, M. and Steurer, A., 1999, *The 'Frontier of Monetisation' in Environmental Valuation*, 6th London Group meeting on environmental accounting, 15–19 November 1999, Canberra, Australia.

OECD, 2011, *Towards Green Growth: Monitoring Progress — OECD Indicators*, OECD, Paris, May 2011, http://www.oecd.org/document/10/0,3746, en_2649_37465_47983690_1_1_1_37465,00.html.

Peskin, H. M, 1972, *National Accounting and the environment*, Artikler 50, Statistiks Sentralbya, Oslo, 1972, http://www.ssb.no/histstat/art/art_050.pdf.

Plan Bleu, 2010, Valeur économique des bénéfices soutenables provenant des écosystèmes marins et côtiers méditerranéens, A. Mangos, J-P. Bassino, D. Sauzade, Plan Bleu, Centre d'Activités Régionales PAM/PNUE, Sophia Antipolis, March 2010.

PNUE/PAM/Plan Bleu, 2004, *L'eau des Méditerranéens : situation et perspectives*. Rapport préparé par Jean Margat et Sébastien Treyer. No. 158, de la Série des rapports techniques du PAM, PNUE/PAM, Athènes, 2004, http://www.planbleu.org/publications/mts158.pdf.

Postel, S. L., Daily, G.C., Ehrlich, P. R., 1996, 'Human Appropriation of Renewable Fresh Water', *Science*, *New Series*, Vol. 271, No. 5250 (Feb. 9, 1996), pp. 785–788.

Rapport D. J., Whitford W. G., 1999, 'How Ecosystems Respond to Stress — Common properties of arid and aquatic systems', in *BioScience* Volume: 49 Number: 3, pp. 193–203, American Institute of Biological Sciences.

Rapport David J., 2007, 'Sustainability science: an ecohealth perspective', in *Integrated Research System for Sustainability Science*, Springer, Sustain Sci (2007) 2:77–84 DOI 10.1007/s11625-006-0016-3.

Ten Brink P. editor, 2011, TEEB D1, TEEB for National and International Policy Makers, An output of TEEB: The Economics of Ecosystems and Biodiversity, Earthscan, http://www.earthscan.co.uk/?tabid=102580 (preliminary draft chapters can be downloaded from http://www.teebweb.org/ForPolicymakers/TEEBforPolicyMakersDraftChapters/tabid/29432/Default.aspx).

The 2030 Water Resource Group, 2009, Charting Our Water Future: Economic Frameworks to Inform Decision-making, the World Bank 2009, http://www.2030waterresourcesgroup.com/water_full/Charting_Our_Water_Future_Final.pdf.

United Nations, 1993, *Integrated Environmental and Economic Accounting* (SEEA 1993), Studies in Method, Series F, No. 61. Department for Economic and Social Information and Policy Analysis, Statistical Division, United Nations, New York, http://unstats.un.org/unsd/envaccounting/pubs.asp.

United Nations, 2009, System of National Accounts 2008, European Communities, International Monetary Fund, Organisation for Economic Co-operation and Development, United Nations and the World Bank, ISBN 978-92-1-161522-7, http://unstats.un.org/unsd/nationalaccount/docs/SNA2008.pdf.

UNSD, 2003, Handbook of National Accounting: Integrated Environmental and Economic Accounting 2003 (SEEA 2003) UNSD, EC, OECD, WB and IMF (2003), http://unstats.un.org/unsd/envaccounting/seea2003.pdf.

UNSD, 2007, System of Environmental-Economic Accounting for Water (SEEAW), United Nations Statistics Division, 2007, http://unstats.un.org/unsd/envaccounting/seeaw/seeawdraftmanual.pdf.

UNSD, 2011, SEEA Experimental Ecosystem Accounts: A Proposed Outline and Road Map, Paper prepared by UNSD, EEA and the World Bank, ESA/STAT/AC.238, UNCEEA/6/6, Sixth Meeting of the UN Committee of Experts on Environmental-Economic Accounting, New York, 15–17 June 2011, http://unstats.un.org/unsd/envaccounting/ceea/meetings/UNCEEA-6-6.pdf.

UNSD, 2011, Economic Environmental Accounting Library — Database Search, http://unstats.un.org/unsd/envaccounting/ceea/archive/searchkey2.asp.

UNESCWA, 2009, Compendium of Environment Statistics in the ESCWA Region, E/ESCWA/SD/2009/13, December 2009, http://www.escwa.un.org/information/publications/edit/upload/sd-09-13.pdf.

UNESCWA, 2010, 'Water Statistics and Water Accounts in Jordan', presentation by Suna Abou Zahra, http://www.escwa.un.org/divisions/events/Countrypapers/JordanDOSPresentation.pdf.

UNESCWA, 2011a, Environment Statistics, Indicators and Accounts Project (ESIAP), http://www.escwa.un.org/esiap/.

UNESCWA, 2011b, Statistical Information System ESIS, http://esis.escwa.org.lb/?id=18.

UNU, 2010, Satoyama-Satoumi, Ecosystems and Human Well-being: Socio-ecological Production Landscapes of Japan – Summary for Decision Makers. United Nations University, Tokyo, Japan, 2010, http://www.ias.unu.edu/resource_centre/SDM-EN_24Feb2011.pdf.

Valero, 2006, Fundamentals of Physical Hydronomics: a new approach to assess the environmental costs of the European Water Framework Directive, Antonio Valero, Javier Uche, Alicia Valero, Amaya Martínez, José Manuel Naredo and Joan Escriu, ISEE 2006, http://www.fnca.eu/fnca/docu/docu230.pdf.

Vanoli, A., 2005, *A History of National Accounts*, IOS press, 2005, ISBN: 1-59603-469-3 http://www.iospress.nl/flyers_b/fl1586034693.pdf.

Weber, J.-L., 2007, 'Implementation of land and ecosystem accounts at the European Environment Agency', *Ecological Economics*, Volume 61, Issue 4, 15 March 2007, p. 695–707, http://www.sciencedirect.com/science/article/pii/S0921800906004629.

Weber, J.-L., 2008, Land & ecosystem accounts in the SEEA revision. Position paper for the London Group meeting, Brussels, 29 September–3 October 2008, http://unstats.un.org/unsd/envaccounting/londongroup/meeting13/LG13_25a.pdf.

Weber J.-L., 2011a, Ecosystem Services in Green National Accounting, Jean-Louis-Weber, in Ecosystem Services and Global Trade of Natural Resources. Thomas Koellner Editor, Routledge, 2011.

Weber, J.-L., 2011b, Approach to Simplified Ecosystem Capital Accounts, Jean-Louis Weber, European Environment Agency, UNCEEA/6/33, Sixth Meeting of the UN Committee of Experts on Environmental-Economic Accounting, New York, 15–17 June 2011, http://unstats.un.org/unsd/envaccounting/ceea/meetings/UNCEEA-6-33.pdf.

Wiedmann, T., 2008, Development of an Embedded Carbon Emissions Indicator – Producing a Time Series of Input-Output Tables and Embedded Carbon Dioxide Emissions for the UK by Using a MRIO Data Optimisation System, Wiedmann, T., Wood, R., Lenzen, M., Minx, J., Guan, D. and Barrett, J., Report to the UK Department for Environment, Food and Rural Affairs by Stockholm Environment Institute at the University of York and Centre for Integrated Sustainability Analysis at the University of Sydney, June 2008, DEFRA, London, United Kingdom, http://www.isa.org.usyd.edu.au/publications/documents/Defra_EmbeddedCarbon_Main.pdf.

World Bank, 1989, Environmental accounting for sustainable development (selected papers from joint UNEP/World Bank workshops from 1983), by Ahmad, Yusuf J., El Serafy, Salah, Lutz, Ernst (editors), the World Bank, Washington, DC, 1989.

World Bank, 2011, *The Changing Wealth of Nations: Measuring Sustainable Development in the New Millennium*, the World Bank, 2011, http://siteresources.worldbank.org/ENVIRONMENT/Resources/ChangingWealthNations.pdf.

World Bank, 2011, Wealth Accounting and the Valuation of Ecosystem Services (WAVES): A Global Partnership, Paper prepared by the World Bank, ESA/STAT/AC.238, UNCEEA/6/7, Sixth Meeting of the UN Committee of Experts on Environmental-Economic Accounting, New York, 15–17 June 2011, http://unstats.un.org/unsd/envaccounting/ceea/meetings/UNCEEA-6-7.pdf.

Annex 1 List of tables and accounting items

List of tables and accounting items as labelled in the mock-up tables of Simplified Ecosystem Capital Accounts (supplied as MS Excel worksheet). Where they are based on real data, the mock-up numbers refer in principle to \sim 1995 and \sim 2005

Headings

Ecosystem classes

	Inland ecosystem landscapes					Sub-		S	ea		Atmo- sphere			
Dominant urban landscape	Dominant agri- culture/ cropland	agri- culture/	nant forested landscape	Other domi- nant natural land- scape	Composite land- scape	Sub- total Land	River systems	total inland eco- systems	Fisheries (EEZ, all fishing areas)	Inter- national fishing areas	Total Fisheries	Regula- tion potential	Regula- tion potential	Grand total

And/or ISIC classes

Rows (part 1): Accounts in physical units

[A] Land-cover stocks and flows basic account

A1 Total EU-27 1990, km²

a11	Artificial	curfocos
all	Artiliciai	Surfaces

- a12 Large to medium field arable land & shrub crops
- a13 Pastures & mosaic farmland (small fields)
- a14 Forests cover
- a15 Natural grassland, shrubs
- a16 Open space with little or no vegetation
- a17 Wetlands
- a18 Water bodies

A2 Land cover change, total flows 1990–2006, km² [lcf1+lcf2+lcf3]

- a21 lf1 Land development processes, urban sprawl, land use intensification
 - a211 If11 Artificial development over agriculture
 - a212 If12 Artificial development over forests
 - a213 If13 Artificial development of other natural land cover
 - a214 If14 Conversion from small fields agriculture and pasture to broad pattern cropland
 - a215 lf15 Conversion from forest to agriculture
 - a216 lf16 Conversion from marginal land to agriculture
 - a217 If17 Water bodies creation and management

a22 lf2 Land restoration processes

- a221 If21 Conversion from crops to set aside, fallow land and pasture
- a222 lf22 Withdrawal of farming
- a223 lf23 Forest creation, afforestation of agriculture land

	a23	lf3 Rota a231 a232 a233 a234 a235	lf31 Internal conversion of artificial surfaces lf32 Internal conversion between agriculture crop types lf33 Recent tree clearing and forest transition lf34 Forests conversions and recruitment lf35 Changes of land cover due to natural and multiple causes				
A3	lf4 No	observe	d land cover change [A1-A2]				
A4	lf5 Cha	nge of d	ominant landscape type [A5-A1]				
A 5	Total E	Total EU-27 2006, km² (as A1)					
	•	-	Carbon/biomass Account: Net Ecosystem Carbon Balance (NECB) & ble Carbon Surplus (NEACS)				
Stock a	accounts						
B1	Stock to b11 b12	Stock t), 10 ⁶ tonnes of C 1 (~1995), 10 ⁶ tonnes of C/Soil 1 (~1995), 10 ⁶ tonnes of C/trees & shrubs				
B2	Stock to b21 b22	Stock t	05), 10 ⁶ tonnes of C 10 (~2005), 10 ⁶ tonnes of C/soil 10 (~2005), 10 ⁶ tonnes of C/trees & shrubs				
В3	Change b31 b32 b33	Change Change	10 ⁶ tonnes of C e t10-t1, 10 ⁶ tonnes of C/soil e t10-t1, 10 ⁶ tonnes of C/trees & shrub nnual C increase %				
B4	Mean a b41 b42 b43 b44 b45 b46 b47 b48	GPP 10 Rp = Re NPP 10 Rh = Re NEP 10 Leakag b46a b46b	arbon/biomass account and NECB 6 tonnes of C espiration by Plants 6 tonnes of C espiration by Heterotrophs and Decomposers 6 tonnes of C espiration by Heterotrophs and Decomposers 7 tonnes of C espiration by Heterotrophs and Decomposers 8 tonnes of C Espiration by Heterotrophs and Decomposers 9 tonnes of C Espiration by Plants 10 tonnes of C Espiration by Heterotrophs and Decomposers 10 tonnes of C Es				
	<i>∪</i> ±0	b481 b482	Net removal/crops b481a total harvest b481b leftovers, returns Net removal/grazing b482a total grazing				
		b483	b482b animal excretion return to pasture Net removal/timber b483a total harvest b483b leftovers, returns				
		b484	Net removal/fish b484a total catches b484b leftovers, returns				
		b485 b486	Removal/extraction of soil, peat Organic fertilisation				

В5

B6

B7

B8

В9

B10

B11

B12

B13

B14

C1

C2

C3

c32b

B8-B7

b49 mean NECB (~1995-~2005), 106 tonnes of C, [b47-b481-b482-b483-b484-b485+b486] mean NECB (~1995-~2005), 106 tonnes of C_soil mean NECB (~1995–~2005), 106 tonnes of C_trees & shrubs b492 **Net Ecosystem Accessible Carbon Surplus** Carbon stress coefficient t1 (~1995) ([b81+b82)/100) A = Total area% WHERE NECB_Soil < or = 0 b51 b52 B = area% of SELU WHERE NECB_Trees & shrubs < NEP surplus Carbon stress coefficient t10 (~2005) A = Total area% WHERE NECB_Soil < or = 0 b62 B = area% of SELU WHERE NECB_Trees & shrubs < NEP surplus Net Ecosystem Accessible Carbon Surplus: NEACS t1 (~1995), weighted 106 tonnes of C [proxy b47*B8] Net Ecosystem Accessible Carbon Surplus: NEACS t10 (~2005), weighted 106 tonnes of C [proxy b47*B9] Change in NEACS Use of biological carbon (removals) t1 (~1995), weighted 106 tonnes of C [b481+b482+b483-b484] Use of biological carbon (removals) t10 (~2005), weighted 106 tonnes of C [b481+b482+b483-b484] Use of fossil carbon, t1 (~1995), 106 tonnes Use of fossil carbon, t1 (~2005), 106 tonnes Ecosystem Accessible Carbon Surplus index t1 (~1995), [B7/B9*100)] [NB should be > 100] Ecosystem Accessible Carbon Surplus index t10 (~2005), [B8/B10*100)] [NB should be > 100] [C] Ecosystem Capital Water Account: Total and Net Ecosystem Accessible Water Surplus (TEAWS and NEAWS) Water stock accounts Water stock t1 (~1995) 106 m3 c11 Aquifers c111 of which aquifers accessible water stock c12 Soil water of which soil accessible water stock c121 Rivers c13 of which rivers accessible water stock c131 c14 Lakes and dams c141 of which lakes and dams accessible water stock Water stock t10 (~ 2005) 10⁶ m³ (as C1) Annual water flows account 106 m³ Precipitation c31 c32 Spontaneous real evapotranspiration

> of which real evapotranspiration induced by rainfed cultivated vegetation of which real evapotranspiration induced by non-cultivated vegetation

- c31-c32 s/t Total available effective rainfall
- c33 Net spontaneous internal and external transfers
- c34 s/t Total available effective rainfall after spontaneous transfers
 - c34a of which Inaccessible runoff (flood...)
 - c34b of which reserved runoff/dillution of pollution, biological needs
 - c34c of which net transfers of pollution as additional reserved runoff/dilution of pollution
 - c34d of which additional Evapotranspiration induced by irrigation and other uses
- c35 Accessible ecosystem water flow [c34-c341-c342-c343-c344]
- c36 Withdrawals of water
 - c361 Withdrawals of fresh water (abstraction, diversion to electricity turbine, net storage in reservoirs)
 - c362 Withdrawals of sea water
- c37 Net transport of water (artificial transfers by mains and canals, conveyance to WWTP...)
- c38 Urban runoff inflow
- c39 Returns of waste water
 - c391 Returns of water/waste water to water bodies incl. urban runoff outflow
 - c392 Returns of water/waste water to the sea
- c40 Returns of water to soil/losses in transport
- c41 Return of water to soil/Irrigation
- c42 Evapotranspiration induced by irrigation and other uses
- c43 Net runoff (external inflows final outflows)

Net Ecosystem Accessible Water Surplus

- C5 Total Ecosystem Accessible Water t1 (~1995) [c12+c14+c16+c18+c35+c37+c391-c38+c40+c41]
- C6 Total Ecosystem Accessible Water t10 (~2005) [c22+c24++c26+c28+c35+c37+c391-c38+c40+c41]
- C6-C5 Change in total accessible water [C6-55]
- C7 Water stress coefficient t1 (~1995), [mean+stdv number of dry days over 30 years/dry days during growing season t1]
- C8 Water stress coefficient t10 (~2005), [mean+stdv number of dry days over 30 years/dry days during growing season t10]
- C9 Net Ecosystem Accessible Fresh Water Surplus t1 (~1995), [C5*(1-C7)] 10⁶ weighted m³
- C10 Net Ecosystem Accessible Fresh Water Surplus t10 (~2005) [C6*(1-C8)], 10⁶ weighted m³
- C10-C9 Change in Net Ecosystem Accessible Fresh Water Surplus [C10-C9]
- C11 Withdrawals of fresh water t1 (~1995) 106 weighted m³
- C12 Withdrawals of fresh water t10 (~2005) 106 weighted m³
- C13 Ecosystem Accessible Water Surplus index t1 (~1995), [((C11-C9)/C9))*100]
- C14 Ecosystem Accessible Water Surplus index t10 (~2005), [((C12-C10)/C10))*100]

[D] Landscape green infrastructure accounts: Landscape Ecosystem Potential (LEP), Green Accessible Landscape Infrastructure (GALI) & Rivers Ecosystem Potential (REP)

Landscape Ecosystem Potential

- D1 Green Background Landscape Index 2000, 5 km smoothing, 10³ points-km², 0–100 scale
 - d11 Mean GBL_P per km²
 - d12 GBLI change 1990–2006
 - d13 Mean change
- D2 Effective Mesh Size index (ln MEFF), 10³ points-km², 0–100 scale
 - d21 Mean MEFF_P per km²
- D3 Stated Social Nature Value index (Naturilis), 10³ points-km², 0–100 scale
 - d31 Mean NAT_P per km²
- D4 Landscape Ecosystem Potential (LEP = f(GBLI, Naturlis, ln MEFF)) t1 (~1995), 10³ points-km², 0–100 scale
 - d41 Mean LEP_P per km²
- D5 Landscape Ecosystem Potential (LEP = f(GBLI, Naturlis, ln MEFF)) t2 (~2005), 10³ points-km², 0–100 scale
 - d51 Mean LEP_P per km²
- D6 Net change in LEP (10³ LEP_P) (D6=D5-D4)
 - d61 Mean annual net change in LEP
 - d62 Mean annual losses in LEP
 - d63 Mean annual gains in LEP
- D7 Mean Landscape Ecosystem Potential (LEP) by km², t1 (~1995), 0–100 scale
- D8 Mean Landscape Ecosystem Potential (LEP) by km², t10 (~2005), 0–100 scale

Green Accessible Landscape Infrastructure

- D9 Green Ecotones Index (GEI)
- D10 Green ecotones index, GEI t1 (~1995), 103 GE_P points
- D11 Green ecotones index, GEI t10 (~ 2005), 10³ GE_P points

D11-D10 Change in GEI

- D12 Mean GEI t1 (~1995)/points by km²
- D13 Mean GEI t10 (~ 2005)/points by km²
- D14 GALI = Green Accessible Landscape Infrastructure Index (SQRT GBLI*GEI), t1 (~1995) d141 Mean GALI per km², t1 (~1995)
- D15 GALI = Green Accessible Landscape Infrastructure Index (SQRT GBLI*GEI), t1 (~2005) d151 Mean GALI per km², t1 (~2005)

Rivers Ecosystem Potential

- D00 River infrastructure in km
- D16 River infrastructure potential in 10³ Standard-River-Kilometer (1 srkm = 1 km*1m³/second)
 - d161 Large rivers
 - d162 Medium rivers
 - d163 Small rivers
 - d164 Brooks, streams
- D17 River integrity composite index, mean value t1 (~1995) [(d171+d172+d173)/3]
 - d171 Water quality
 - d172 Fragmentation
 - d173 Rivers green ecotones
- D18 River integrity composite index, mean value t10 (~2005) [(d181+d182+d183)/3]
 - d181 Water quality
 - d182 Fragmentation
 - d183 Rivers green ecotones
- D19 Rivers Ecosystem Potential (REP) t1 (~1995), weighted 10³ srkm
- D20 Rivers Ecosystem Potential (REP) t10 (~2005), weighted 10³ srkm
- D16 = D20-D19 Change in REP
- D21 Mean Rivers Ecosystem Potential (REP) t1 (~1995)/points by km²
- D22 Mean Rivers Ecosystem Potential (REP) t10 (~2005)/points by km²

[E] Ecosystem Capital Biodiversity Account: Biodiversity Infrastructure Integrity (BII) & Ecosystem's Biodiversity Rating (EBR)

Biodiversity Infrastructure Integrity Index

- E1 BII = GEI weighted LEP & GEI weighted REP, t1 (~1995) [(SQRT D4*D10)] & [(SQRT D19*D10)]
- E2 BII = GEI weighted LEP & GEI weighted REP, t10 (~2005) [(SQRT D6*D11) & [(SQRT D20*D11)]
- E2-E3 Change in BIII
- E2-3 % Change in BIII %
- E4 Mean Biodiversity Infrastructure Integrity index (BII) by km², t1 (~1995)
- E5 Mean Biodiversity Infrastructure Integrity index (BII) by km², t10 (~2005)

Species/biotopes diagnosis

- E5 Species/biotopes diagnosis index, SBD t1 (~1995), 0–100
- E6 Species/biotopes diagnosis index, SBD t10 (~2005), 0–100
- E6-E5 Change in species/biotopes diagnosis index

Ecosystem's Biodiversity Rating

- E11 Mean Ecosystem Biodiversity Rating (EBR) t1 (~1995), weighted km² [SQRT E4*E6]
- E12 Mean Ecosystem Biodiversity Rating (EBR) t10 (~2005), weighted km² [SQRT E5*E7]

[F1] Total Ecosystem Potential Account

Total Ecosystem Potential (TEP) & Net Change, Territorial Ecosystem Capital Degradation (TECD)

Gross Inland, Sea and Atmosphere Ecosystem Potential (NEACS & REP)

- F1 = B7+D19 Gross Inland, Sea and Atmosphere Ecosystem Potential (NEACS & REP) t1 (~1995)
- F2 = B8+D20 Gross Inland, Sea and Atmosphere Ecosystem Potential (NEACS & REP) t10 (~2005)
- (F2-F1)/10 Mean net annual change in NEACS_REP [(B8D11-B7D10)/10]

Limiting factors to C access: access to other services and maintenance of ecosystem functions

- C13 Ecosystem Accessible Water Surplus (EAWS) index t1 (~1995), [((C11-C9)/C9))*100]
- D7 Mean Landscape Ecosystem Potential (LEP) by km², t1 (~1995), 0–100 scale
- D21 Mean Rivers Ecosystem Potential (REP) t1 (~1995)/points by km²
- E11 Mean Ecosystem Biodiversity Rating (EBR) t1 (~1995), weighted km² [SQRT E4*E6]
- C14 Ecosystem Accessible Water Surplus (EAWS) index t10 (~2005), [((C12-C10)/C10))*100]
- D8 Mean Landscape Ecosystem Potential (LEP) by km², t10 (~2005), 0–100 scale
- D22 Mean Rivers Ecosystem Potential (REP) t10 (~2005)/points by km²
- E12 Mean Ecosystem Biodiversity Rating (EBR) t10 (~2005), weighted km² [SQRT E5*E7]
- F3 Mean limiting factors index t1 [(C13+D7+E11)/3] & [(D21+E11)/2]
- F4 Mean limiting factors index t10 [(C14+D8+E12)/3] & [(D22+E12)/2]
- F4-F3 Relative change % = functional gain (+) or loss (-), 0 to 100 scale

Net total ecosystem capital potential [NTECP] & Ecosystem capital degradation [ECD], in Ecosystem Potential Unit Equivalents [1 EPUE = 1 NEACS Unit * functional coefficient]

- F5 Net Total Ecosystem Capital Potential t1 (~1995), in 10³ EPUE [bottomline F5 = F1]
- F6 Net Total Ecosystem Capital Potential t10 (~2005), in 10³ EPUE [F6= (F2*(1-((F4-F3)/F3))]
- F6-F5 NTECP Change in EPUE (-) or (+), period t1t10 (~1995-~2005), in 10³ EPUE
- F6-F5 annual Mean Annual NTECP Change in EPUE (-) or (+), period t1t10 (~1995-~2005), in 103 EPUE
- F6-F5% Mean Annual NTECP Change in EPUE (-) or (+), period t1t10 (~1995-~2005)

- F7 Ecosystem improvement, period t1t10 (~1995–~2005), in 10³ EPUE
 - f71 Effect of Ecosystem restoration programme, mean annual amount period t1t10 (~1995–~2005), in 10³ EPUE
 - f72 Ecosystem spontaneous natural improvement, mean annual amount period t1t10 (~1995—~2005), in 10³ EPUE
- F8 Ecosystem degradation & natural disturbance, 10 years period t1t10 (~1995–~2005), in 10³ EPUE
 - f81 Ecosystem degradation & natural disturbance, mean annual amount, period t1t10 (~1995~~2005), in 10³ EPUE
 - f82 Effect of natural disturbances, mean annual amount period t1t10 (~1995–~2005), in 10³ EPUE
- F9 Ecosystem Capital Degradation (ECD), mean annual amount ~1995—~2005, in 10³ EPUE [F8-F10]
- F9% Mean annual ECD/TEP %, period t1t10 (~1995-~2005)

[F2] Account of territorial ecosystem capital degradation (TECD) by stress factors (in EPUE)

- F9 Territorial Ecosystem Capital Degradation (TECD)
 - f91 Effect of land cover change
 - f911 Urban and infrastructures development over agriculture
 - f912 Conversion of pasture/grassland to cropland
 - f913 Deforestation (forest land uptake by agriculture or urban sprawl)
 - f914 Other shift to more artificial or intensive land cover type
 - f92 Restructuring/destructuring of landscapes and rivers
 - f93 Overexploitation of biological resources
 - f931 Agriculture overharvesting and over grazing
 - f932 Clearing of forest beyond mean NEP
 - f933 Overfishing
 - f934 Overhunting
 - f94 Waste disposal, pollution
 - f941 Pollution/Use of chemicals in agriculture, forestry
 - f942 Pollution/Waste dumping
 - f943 Water pollution
 - f944 Air pollution
 - f945 Emission of GHGs

[G] Demand and Accessibility to Ecosystem Services: Ecosystem Carbon/Biomass, Ecosystem Fresh Water, Green Infrastructure Neighbourhood Ecosystem Services (GINES)

- A1 Total EU-27 1990, km²
 - a11 1 Artificial surfaces, urban land cover EU-27, 1990 (~1995), km² a111 Mean C1 per km², ~1995, %
- A3 Total EU-27 2006, km²
 - a31 1 Artificial surfaces, urban land cover EU-27, 2006 (~2005), km² a311 Mean C1 per km², ~2005, %
- G1 Population 2000 (source: Eurostat+Pop_to_CLC_v5)
 - g11 Population 1995 estimated at 0.98 of 2000
 - g12 Population 2005 estimated at 1.02 of 2000

G2 Net Accessible Ecosystem Carbon Net Accessible Ecosystem Carbon per capita 1995 (tonnes) [g21 = B7/g11]g21 g22 Net Accessible Ecosystem Carbon per capita 2005 (tonnes) [g22 =B8/g11] G3 Net Accessible Ecosystem Fresh Water Net Ecosystem Accessible Fresh Water per capita 1995 (m³) [g31 = C9/g11] g31 g32 Net Ecosystem Accessible Fresh Water per capita 2005 (m³) [g32 = C10/g12] G4 Accessible landscape services/Green Infrastructure Neighbourhood Ecosystem Services (GINES) G41= D14 GALI = Green Accessible Landscape Infrastructure Index (SQRT GBLI*GEI), t1 (~1995) g411= d141 Mean GALI per km², t1 (~1995) G42= D15 GALI = Green Accessible Infrastructure Landscape Index (SQRT GBLI*GEI), t1 (~2005) g421= d151 Mean GALI per km², t1 (~2005) G43 Demand of GINES_5km (= SQRT GALI * a11 Artificial) ~1995 Mean GINES 5km demand per km² ALL g431 g432 Mean GINES 5km demand per km² C1 Artificial G44 Demand of GINES_5km (= SQRT GALI * a11 Artificial) ~2005 g441 Mean GINES_5km demand per km² ALL g442 Mean GINES_5km demand per km² C1 Artificial G45 Mean accessibility of GINES_5km (GALI/a11 Artificial) t1 ~1995 G46 Mean accessibility of GINES_5km (GALI/a31 Artificial) t1 ~2005 G47 Accessible GINES/landscape services (G47 = (G43*G45)*g11), 106 points G48 Accessible GINES/landscape services (G48 = (G44*G46)*g12), 10⁶ points [H] Physical Balance Sheet: Assets and Liabilities Physical Assets [in EPUE] H1=F5 Opening Balance Sheet: Net Total Ecosystem Capital Potential, in 10³ EPUE [bottomline F5 = F1] H11=F5 Non-financial ecosystem assets H111 Land ecosystems H112 River ecosystems H113 H114 Atmosphere H12 Financial ecosystem assets (in 10³ EPUE)

Change in Total Ecosystem Potential & Ecosystem capital degradation

- H2 Change in Ecosystem Potential Due to Economic Activities f71 Effect of Ecosystem restoration programme, in 10³ EPUE F9 Territorial Ecosystem Capital Degradation (TECD), in 10³ EPUE [F9=F8-F10] Н3 Other Change in Potential of Ecosystem Capital f72 Ecosystem spontaneous natural improvement, in 10³ EPUE f82 Effect of natural disturbances, in 10³ EPUE H4 Net Change in Physical Ecosystem Assets NTECP (-) or (+) [L5 = f71+f72-F9-F8] H41 Net change in non-financial ecosystem assets h411 Land ecosystems h412 River ecosystems h413 Sea h414 Atmosphere H42 Net acquisition of new ecosystem physical assets (Ecosystem improvement, ECD embedded into exports)
- H5=F6 Closing Balance Sheet: Total Ecosystem Potential, in 10³ EPUE [H5=(F2*(1-((F4-F3)/F3))] (detail as H1)

Physical Liabilities [in EPUE]

- H6 Opening Balance Sheet Total Ecosystem Potential, in 10³ EPUE [bottomline F5 = F1]
- H7 Ecosystem restoration targets (recovery from historical damages, compliance to conventions/ regulations)
 - h71 National targets
 - h72 International targets
 - h73 Private targets
 - h74 Change in ecosystem restoration targets
- H8 Acquisition of new physical liabilities
 - h81=F9 Territorial Ecosystem Capital Degradation (TECD), in 103 EPUE [F9=F8-F10] of t-1
- K2 Ecosystem capital degradation in 'consumed imports', agriculture & forest, in EPUE
- K3 Ecosystem capital degradation in 'consumed imports', fisheries, in EPUE
- K4 Ecosystem capital degradation in 'consumed imports', atmosphere CO₂-e potential, in EPUE
- H9 Reduction of physical liabilities
 - h91=f71 Reduction of physical liabilities by ecosystem restoration programmes
 - h92=f72 Ecosystem spontaneous natural improvement, in 103 EPUE
 - h93 Reduction of physical liabilities by acquisition of EPUE (mitigation/compensation)
 - h94 Reduction of physical liabilities by swaps and debts consolidation
- H10 Net change in physical liabilities (=h74+H8-H9)
- H11 Closing Balance Sheet, in 10³ EPUE

Rows (part 2): Accounts in money

[I] Estimation of unit costs of ecosystem capital restoration by stress factors

f91 & j11 Effect of land cover change

- f911 & j111 Urban and infrastructures development over agriculture ==> compensation
- f912 & j112 Conversion of pasture/grassland to cropland => set aside, loss of crop revenue
- f913 & j113 Deforestation (forest land uptake by agriculture or urban sprawl) => reforestation
- f914 & j114 Other shift to more artificial or intensive land cover type => compensation

f92 & j12 Restructuring/destructuring of landscapes and rivers ==> plantation of hedgerows

f93 & j13 Overexploitation of biological resources

f931 & j131	Agriculture overharvesting and over grazing ==> yield abatement, organic
	fertilisation, change of crop

f932 & j132 Clearing of forest beyond mean NEP ==> yield abatement

f933 & j133 Overfishing ==> yield abatement

f934 & j134 Overhunting

f94 & f14	Waste disposal, pollution ==> yield abatement					
	f941 & j141	Pollution/Use of chemicals in agriculture, forestry ==> yield abatement less				
		cost of chemicals				
	f943 & j142	Water pollution ==> cost of abatement programmes				
	f942 & j143	Pollution/Waste dumping ==> cost of restoration programmes				
	f944 & j144	Air pollution ==> cost of abatement programmes				
	f945 & j145	Emission of GHGs ==> investments in clean technologies				

[J] Ecosystem Capital Depreciation: Estimation of Territorial Consumption of Ecosystem Capital in 10⁶ Euro

J1	Territo	orial Consumption of Ecosystem Capital, mean annual value period t1t10 in 106 Euro				
	j11	Effect of land cover change				
		j111 Urban and infrastructures development over agriculture				
		j112 Conversion of pasture/grassland to cropland				
		j113 Deforestation (forest land uptake by agriculture or urban sprawl)				
		j114 Other shift to more artificial or intensive land cover type				
	j12	Restructuring/destructuring of landscapes and rivers				
	j13	Overexploitation of biological resources				
	,	j131 Agriculture overharvesting and over grazing				
		j132 Clearing of forest beyond mean NEP				
		j133 Overfishing				
		j134 Overhunting				
	f14	Waste disposal, pollution				
		j141 Pollution/Use of chemicals in agriculture, forestry				
		j142 Water pollution				

[K] Account of Ecosystem Capital Degradation & Depreciation Embedded into Imports & Exports, in EPUE & 10^6 Euro

K1	Virtual or embedded land in 'consumed imports', agriculture, km ²
K2	Ecosystem capital degradation in 'consumed imports', agriculture & forest, in EPUE
K3	Ecosystem capital degradation in 'consumed imports', fisheries, in EPUE
K4	Ecosystem capital degradation in 'consumed imports', atmosphere CO ₂ -e potential, in EPUE
K5	Unpaid ecosystem depreciation/'consumed imports', agriculture & forest, at EU mean price
K6	Unpaid ecosystem depreciation/'consumed imports', fisheries potential
K7	Unpaid ecosystem depreciation/'consumed imports', CO ₂ -e potential
K8	Ecosystem capital depreciation virtually embedded into imports (total)
K9	Ecosystem capital depreciation virtually embedded into exports (total)

[L] Sustainable Ecosystem Services Macro-economic Benefits:

Primary production, basic price

Pollution/Waste dumping

Air pollution Emission of GHGs

j143 j144

j145

Ecologically Sustainable Total Induced Value Added (ES-TIVA), in 106 Euro (by sectors/ISIC)

L2	Value added of primary production
L3	Subsidies to primary production
L4	Ecosystem capital degradation resulting from economic exploitation %
L5	Total value added induced by primary production of agriculture product
L6	Ecologically Sustainable TIVA/ (sustainable TIVA)/agriculture products
L7	Total value added induced by primary production of forestry products
L8	Ecologically Sustainable TIVA/ (sustainable TIVA)/forestry products

L1

value added induced by primary production of fishing products
ogically Sustainable TIVA/ (sustainable TIVA)/fishing products
value added induced by primary production of fresh water supply
ogically Sustainable TIVA/ (sustainable TIVA)/water supply
)

[M] Economic aggregates and additional adjustments for CEC, 106 current Euro, EU-27

M01 GDP
M02 Final Consumption
M03 Imports CIF
M04 Exports FOB
M05 Consumption of Fixed Capital

Consumption of Ecosystem Capital

M1	Territorial Consumption of Ecosystem Capital (TCEC) (M1=J1)				
	m11=J1/land	Territorial Consumption of Ecosystem Capital, in 10 ⁶ Euro — Inland ecosystems			
	m12=J1/fish.	Territorial Consumption of Ecosystem Capital, in 10 ⁶ Euro — Sea/fisheries			
	m13=J1/clim	Territorial Consumption of Ecosystem Capital, 10 ⁶ Euro — Atmosphere/climate			
K8	Ecosystem cap	osystem capital depreciation virtually embedded into imports (total)			
M2	2 Gross Domestic Consumption of Ecosystem Capital (GDCEC) (M2=M1+L8)				

M3 Net Domestic Consumption of Ecosystem Capital (M2=M1+L8-L9)

Adjusted national accounts aggregates

M06 (Conventional) Net Domestic Product (M06=M01-M05)
 m06% % Conventional Net Domestic Product/GDP
 M4 GDCEC Adjusted Net Domestic Product (M4=M01+M2)
 M4% % GDCEC Adjusted Net Domestic Product/GDP
 M5 Final Consumption at Full Price (M5=M02+M3)
 M5% % Final Consumption at Full Price/FC Purchaser Price
 K10 Imports at Full Price
 K10 % Imports at Full Price/Imports CI
 K11 Export at Full Price
 K11 % Exports at Full Price/Export FOB

[N] Monetary Balance Sheet: Assets and Liabilities

Monetary Assets [in 10⁶ Euro]

N1	Ecosys n11 n12 n13	tem Potential Opening Balance Sheet, in 10 Euro — Non Relevant (NR) Market value of ecosystem economic non-financial assets, 10 ⁶ Euro (from SEEA vol.1) Financial ecosystem assets, 10 ⁶ Euro Market value of ecosystem public good assets, in 10 ⁶ Euro — Non Relevant (NR)
N2	Chang n21	e in Ecosystem Potential Due to Economic Activities (+) Effect of ecosystem restoration programmes, in 10 ⁶ Euro (N21= f71 in EPUE*Unit price) n211 Effect of Ecosystem restoration programmes, in 10 ⁶ Euro — Inland ecosystems n212 Effect of Ecosystem restoration programmes, in 10 ⁶ Euro — Fisheries n213 Effect of Ecosystem restoration programmes, in 10 ⁶ Euro — Atmosphere/climate
	n22	(-) Territorial Consumption of Ecosystem Capital (TCEC) (n71=M1=J1) n221 Territorial Consumption of Ecosystem Capital, in 10 ⁶ Euro — Inland ecosystems n222 Territorial Consumption of Ecosystem Capital, in 10 ⁶ Euro — Sea/fisheries n223 Territorial Consumption of Ecosystem Capital, 10 ⁶ Euro — Atmosphere/climate

N3 Counterpart of Other Change in Volume of Ecosystem Capital (+) Ecosystem spontaneous natural improvement, in 106 Euro (n31= f72 in EPUE*Unit price) n31 n32 (–) Effect of natural disturbances, in 10⁶ Euro (n32= f82 in EPUE*Unit price) N4 Net Monetary Change in Ecosystem Assets (–) or (+) Net Monetary Change in non-financial Ecosystem Assets [N4=n21+n31-n22-n32] n411 Land ecosystems n412 River ecosystems n413 Sea n414 Atmosphere n42 Net Acquisition of New ecosystem Financial Assets N₅ Ecosystem Potential Closing Balance Sheet, in 10 Euro — Non Relevant (NR) (detail as N1) Financial Liabilities [in 106 Euro] N6 Opening Balance Sheet Distance to ecosystem restoration targets (historical damages, conventions/regulations) National targets/cost of programmes in 10⁶ Euro n612 International targets/cost of programmes in 106 Euro n613 Private targets/cost of programmes in 106 Euro n64 Change in ecosystem restoration targets n65 Revaluation of programmes cost N7 Acquisition of New Other Financial Liabilities (+) Territorial Consumption of Ecosystem Capital (TCEC) (n71=M1=J1) n711 Territorial Consumption of Ecosystem Capital, in 10⁶ Euro — Inland ecosystems n712 Territorial Consumption of Ecosystem Capital, in 10⁶ Euro — Sea/fisheries n713 Territorial Consumption of Ecosystem Capital, 10⁶ Euro — Atmosphere/climate n72 (+) Ecosystem capital depreciation virtually embedded into imports (total) Ecosystem capital depreciation embedded in 'consumed imports', agriculture & forest, in EPUE n722 Ecosystem capital depreciation embedded in 'consumed imports', fisheries, in EPUE n723 Ecosystem capital depreciation embedded in 'consumed imports', atmosphere CO₂-e potential, in EPUE H8 Reduction of Financial liabilities h81=n21 (-) Reduction of financial liabilities by ecosystem restoration programmes, in 106 Euro (N21= f71 in EPUE*Unit price) Effect of Ecosystem restoration programmes, in 10⁶ Euro — Inland ecosystems h812 Effect of Ecosystem restoration programmes, in 10⁶ Euro — Fisheries h813 Effect of Ecosystem restoration programmes, in 10⁶ Euro — Atmosphere/climate

h82=n31 (-) Ecosystem spontaneous natural improvement, in 106 Euro (n31 = f72 in EPUE*Unit price)

(-) Reduction of Financial liabilities by swaps and debts consolidation

(-) Reduction of Financial liabilities by acquisition of EPUE (mitigation/compensation)

- H9 Net change in Financial liabilities (= h64+H7-H8)
- H10 Closing Balance Sheet

h83

h84

Annex 2 Mock-up accounts

The mock-up accounts can be downloaded from

http://eea.eionet.europa.eu/Public/irc/eionet-circle/leac/library?l=/cube/ecosystem_23sept2011xls/_ $EN_1.0_&a=d.$

43

European Environment Agency

An experimental framework for ecosystem capital accounting in Europe

 $2011 - 43 \text{ pp.} - 21 \times 29.7 \text{ cm}$

ISBN 978-92-9213-233-0 ISSN 1725-2237 doi:10.2800/83749

HOW TO OBTAIN EU PUBLICATIONS

Free publications:

- via EU Bookshop (http://bookshop.europa.eu);
- at the European Union's representations or delegations. You can obtain their contact details on the Internet (http://ec.europa.eu) or by sending a fax to +352 2929-42758.

Priced publications:

• via EU Bookshop (http://bookshop.europa.eu).

Priced subscriptions (e.g. annual series of the *Official Journal of the European Union* and reports of cases before the Court of Justice of the European Union):

• via one of the sales agents of the Publications Office of the European Union (http://publications.europa.eu/others/agents/index_en.htm).

European Environment Agency Kongens Nytorv 6 1050 Copenhagen K Denmark

Tel.: +45 33 36 71 00 Fax: +45 33 36 71 99

Web: eea.europa.eu Enquiries: eea.europa.eu/enquiries





