Category		Title	
GG		General guidance	
		Key category analysis and methodological choice	
Version Guidebook 2009			

Lead author Justin Goodwin

Contents

1	Intr	oduction 3				
	1.1	Categories	3			
	1.2	Fuels	3			
	1.3 Method tiers					
	1.4	Key categories	5			
2	Key	/ category analysis	5			
	2.1	Purpose of key category analysis	5			
	2.1 Fulpose of key category analysis2.2 General approach to identify key categories					
	2.3	General rules for identification of key categories	7			
	2.4	Methodological approaches to identify key categories	12			
	2.5	Reporting and documentation	21			
	2.6	Examples of key category analysis	21			
3						
4	Poi	nt of enquiry	.26			
A		ix A Combining air pollutant specific key categories				

1 Introduction

This chapter explains the tiered approach used to define the generic levels of detail and rigour required for inventory compilation and how to identify the key categories (¹) for which higher tiered methods are required. This chapter has been extensively adapted from the 2006 IPCC (Intergovernmental Panel on Climate Change) Guidelines for National Greenhouse Gas Inventories (IPCC Guidelines) to facilitate estimating transboundary air pollutant emissions.

Methodological choice for individual source categories is important in managing overall inventory quality and minimising uncertainty. Generally, inventory uncertainty is lower when emissions are estimated using the most rigorous, higher tiered, methods (see subsection 1.3) provided for each category or subcategory in the sectoral volumes of this Guidebook. However, these methods generally require more extensive resources for data collection and calculation, so it may not be feasible to use most rigorous method for every category of emissions. It is, therefore, good practice to identify and prioritise the effort on those categories which make the greatest contribution to the overall inventory estimates (and where possible, the uncertainty).

1.1 Categories

The categories used are those of the Nomenclature for Reporting (NFR) as used for reporting emissions according to the United Nations Economic Commission for Europe (UNECE) Guidelines for Reporting Emission Data under the Convention on Long-Range Transboundary Air Pollution (hereinafter the LRTAP Reporting Guidelines) (UNECE 2009)

1.2 Fuels

Fuels are those based on the definitions of the International Energy Agency (IEA) and used in the IPCC Guidelines.

1.3 Method tiers

Emissions can be estimated at different levels of complexity. Within the IPCC Guidelines and adopted by this Guidebook, these are expressed in three tiers of increasing complexity.

The 'Tier 1' method is a 'simple' method using default emission factors only. To upgrade a Tier 1 to a Tier 2 method, the default emission factors should be replaced by country-specific or technology-specific emission factors. This might also require a further split of the activity data over a range of different technologies, implicitly aggregated in the Tier 1 method. A Tier 3 method could be regarded as a method that uses the latest scientific knowledge in more sophisticated approaches and models; more detailed definitions follow.

Tier 1: A method using readily available statistical data on the intensity of processes (activity rates) and default emission factors. These emission factors assume a linear relation between the intensity of the process and the resulting emissions. The Tier 1 default emission factors also assume an average or typical process description. This method is the simplest method, has the highest level of uncertainty and should not

3

^{(&}lt;sup>1</sup>) In Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (GPG2000, IPCC, 2000), the concept was named 'key source categories' and dealt with the inventory excluding the Land Use, Land-Use Change and Forestry (LULUCF) Sector. This definition is elaborated for the purposes of this Guidebook in subsection 1.4.

be used to estimate emissions from key categories (see subsection 1.4).

- Tier 2:Tier 2 is similar to Tier 1 but uses more specific emission factors developed on the
basis of knowledge of the types of processes and specific process conditions that
apply in the country for which the inventory is being developed. Tier 2 methods are
more complex, will reduce the level of uncertainty, and are considered adequate for
estimating emissions for key categories.
- Tier 3: Tier 3 is defined as any methodology more detailed than Tier 2; hence there is a wide range of Tier 3 methodologies. At one end of the range there are methodologies similar to Tier 2 (i.e. activity data x emission factor) but with a greater disaggregation of activity data and emission factors. At the other end of the range are complex, dynamic models in which the processes leading to emissions are described in great detail.

The key criterion to be met before a Tier 3 methodology can replace a Tier 2 methodology is a more accurate estimation of the relevant emissions, reducing the following common sources of error.

- Model error: the extent to which the mathematical representation of the processes underlying the emissions deviate from reality.
- Parameter error: the error in the model parameters (e.g. emission factors, coefficients, etc.)
- Input error: the error in activity data.
- Process error: error introduced through mistakes in the process of compiling the inventory. For complex models, this includes errors in the software implementation of the model.

For small extensions to Tier 2 methodologies, such as the inclusion of abatement measures or refinements to emission factors, it is sufficient to document the quality assurance/quality control (QA/QC) process by which the revised/additional emission factors and associated activity data were obtained. However, where parties wish to use complex simulation models in inventory construction, the model is quite likely to have been developed by a third party. If the use of such models within a methodology is to be accepted as Tier 3, it is necessary to ensure that QA/QC criteria are met by the complex model, the process of parameterisation, and the input data necessary to run the model. These criteria must acknowledge that reviewers should be able to review the methodology within a reasonable time period and are listed below.

- For the inventory construction process to be sufficiently transparent, the model documentation has to be clear, correct, concise, comprehensible, and consistent.
- The scientific quality of the model has to have been documented in peerreviewed publications.
- The model has to have been tested successfully in the situations for which it will be used in inventory construction, implying that model parameters are available for those situations.

• The input data required by the model must be available, and of adequate quality, at the spatial and temporal scales for which the model is to be used.

These criteria are valid for all Tier 3 methodologies. However, they may require further interpretation for sector-specific applications and additional criteria may also be appropriate. For example, where it is appropriate, models intended to be used as Tier 3 methods should demonstrate that they obey the law of conservation of matter.

1.4 Key categories

A key category is one that is prioritised within the national inventory system because it is significantly important for one or a number of air pollutants in a country's national inventory of air pollutants in terms of the absolute level, the trend, or the uncertainty in emissions (as defined in subsection 2.4).

It is good practice for each country to use key category analysis systematically and objectively as a basis for choosing methods of emission calculation. Such a process will lead to improved inventory quality as well as greater confidence in the resulting estimates. The approach for key category analysis is presented in Section 2.

2 Key category analysis

2.1 Purpose of key category analysis

It is good practice for each country to identify its national key categories in a systematic and objective manner. This can be achieved by a quantitative analysis of the relationship between the magnitude of emission in any one year (level) and the change in emission year to year (trend) of each category's emissions compared to the total national emissions.

It is also good practice to focus the available resources for improvement in data and methods on categories identified as *key*. The identification of key categories in national inventories enables the limited resources available for preparing inventories to be prioritised; more detailed, higher tier methods can then be selected for key categories. Inventory compilers should use the category-specific methods presented in sectoral decision trees in the sectoral volumes.

For many sources, higher tier (Tier 2 and 3) methods are usually suggested for key categories. For guidance on the specific application of this principle to key categories, it is good practice to refer to the sector-specific decision trees and guidance for the respective category in the sectoral volumes. In some cases, inventory compilers may be unable to adopt a higher tier method due to lack of resources. This may mean that they are unable to collect the required data for a higher tier or are unable to determine country-specific emission factors and other data needed for Tier 2 and 3 methods. In these cases, although this is not accommodated in the category-specific decision trees, a Tier 1 approach can be used, and this possibility is identified in Figure 2-1. It should, in these cases, be clearly documented why the methodological choice was not in line with the sectoral decision tree. Any key categories where the good practice to pay additional attention to key categories with respect to QA/QC as described in Chapter 6 'Inventory management, improvement and QA/QC', and the separate 'Verification' and 'Inventory QA/QC' sections of each sectoral chapter.




Figure 2-1 Decision tree to choose a good practice method

2.2 General approach to identify key categories

Any inventory compiler should be able to identify key categories in a basic single-year inventory in terms of their contribution to the absolute level of national emissions. For those inventory compilers who have prepared a time series, the quantitative determination of key categories should also include an evaluation of both the absolute level and the trend of emissions. Some key categories may be identified only when their influence on the trend of the national inventory is taken into account.

Subsection 2.3 sets out general rules for the identification of key categories including the sectoral level into which the categories should be split. The methodology follows the principles developed by the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (Volume 1, Chapter 4), described in subsection 2.4, to generate a key category list for each pollutant. However, where a single prioritised list of key categories is needed (e.g. to help in prioritising emission inventory improvements), a further, optional step (Step 2) can be used to consolidate the key categories (Appendix A).

2.3 General rules for identification of key categories

The results of the key category identification will be most useful if the analysis is done at the appropriate disaggregation level of categories. Table 2-1 lists the recommended source categories and, where relevant, identifies special considerations related to the disaggregation of the analysis. For example, the combustion of fossil fuels is a large emission source category that can be broken down into subcategories of first, second or third order, and even to the level of individual plants or boilers. Countries may adapt the recommended level of analysis in Table 2-1 to their national circumstances. In particular, countries using Approach 2 (subsection 2.4) will probably choose the same level of aggregation used for the uncertainty analysis. In some cases, disaggregation to very low levels should be avoided since it may split an important aggregated category into many small subcategories that are no longer *key*. The following guidance describes good practice in determining the appropriate level of disaggregation of categories to identify key categories.

- The analysis should be performed at the level of NFR categories or subcategories at which the Guidebook methods and decision trees are provided in the sectoral volumes. Where possible, and as indicated in Table 2-1, some categories should be disaggregated by main fuel types.
- Each air pollutant emitted from each category should be considered separately. For example, carbon monoxide (CO), nitrogen oxides (NO_x) and particulate matter (PM₁₀) are emitted from each of the detailed road transport classes. The key category analysis for this source should be performed for each of these separately as the methods, emission factors and related uncertainties will be different. In contrast, a collective analysis of all chemical species of non-methane volatile organic compounds (NMVOCs) is appropriate for this category.
- Countries may choose to perform the quantitative analysis at a more disaggregated level than suggested in Figure 2-1 (²). In this case, possible cross-correlations between categories and/or subcategories should be taken into account when performing the key category analysis. When using Approach 2, the assumptions about such correlations should be the same as when assessing uncertainties and identifying key categories (see Chapter 5, Uncertainties).
- The categories included in Table 2-1 below are those for which estimation methods are provided in the sectoral volumes. If countries develop estimates for new categories or gases, these should be added to the analysis under Miscellaneous for the appropriate sector.

7

 $[\]binom{2}{2}$ Most correlations between categories can be avoided by using the aggregation level of this table. Some correlations remain, e.g. in fuel use between stationary combustion and processes. In practice, the effect of correlations for key category analysis should be taken into account in the disaggregation level used for the Approach 2 assessment (for more advice on correlations in uncertainty analysis, see Chapter 3, Data Collection.)

For each key category (see Table 2-1 below), the inventory compiler should determine if certain subcategories are particularly significant. Usually, for this purpose, the subcategories should be ranked according to their contribution to the aggregate key category. Those subcategories that contribute together more than 60 % to the key category should be treated as particularly significant. It may be appropriate to focus efforts towards methodological improvements of these most significant subcategories.

Source	categories to be assessed in key category analysis	Special considerations		
Category code ^b	Category title ^b			
1.A.1.a	Public electricity and heat production	Disaggregate to main fuel types		
1.A.1.b	Petroleum refining	Disaggregate to main fuel types		
1.A.1.c	Manufacture of solid fuels and other energy industries	Disaggregate to main fuel types		
1.A.2.a	Stationary combustion in manufacturing industries and construction: iron and steel	Disaggregate to main fuel types		
1.A.2.b	Stationary combustion in manufacturing industries and construction: non-ferrous metals	Disaggregate to main fuel types		
1.A.2.c	Stationary combustion in manufacturing industries and construction: chemicals	Disaggregate to main fuel types		
1.A.2.d	Stationary combustion in manufacturing industries and construction: pulp, paper and print	Disaggregate to main fuel types		
1.A.2.e	Stationary combustion in manufacturing industries and construction: food processing, beverages and tobacco	Disaggregate to main fuel types		
1.A.2.f.i	Stationary combustion in manufacturing industries and construction: other	Disaggregate to main fuel types		
1.A.2.f.ii	Mobile combustion in manufacturing industries and construction	Disaggregate to main fuel types		
1.A.3.a.ii.(i)	Civil aviation (domestic, landing/take-off (LTO))			
1.A.3.a.ii.(ii)	Civil aviation (domestic, cruise)			
1.A.3.b.i	Road transport: passenger cars			
1.A.3.b.ii	Road transport: light-duty vehicles			
1.A.3.b.iii	Road transport: heavy-duty vehicles			
1.A.3.b.iv	Road transport: mopeds & motorcycles			
1.A.3.b.v	Road transport: gasoline evaporation			
1.A.3.b.vi	Road transport: automobile tyre and brake wear			
1.A.3.b.vii	Road transport: automobile road abrasion			
1.A.3.c	Railways			
1.A.3.d.ii	National navigation (shipping)	Disaggregate to main fuel types		
1.A.3.e	Pipeline compressors	Disaggregate to main fuel types		
1.A.4.a.i	Commercial/institutional: stationary	Disaggregate to main fuel types		

 Table 2-1
 Suggested aggregation level of analysis for Approach 1^a

8

Sourc	e categories to be assessed in key category analysis	Special considerations		
Category code ^b	Category title ^b			
1.A.4.a.ii	Commercial/institutional: mobile	Disaggregate to main fuel types		
1.A.4.b.i	Residential: stationary plants	Disaggregate to main fuel types		
1.A.4.b.ii	Residential: household and gardening (mobile)	Disaggregate to main fuel types		
1.A.4.c.i	Agriculture/forestry/fishing: stationary	Disaggregate to main fuel types		
1.A.4.c.ii	Agriculture/forestry/fishing: off-road vehicles and other machinery	Disaggregate to main fuel types		
1.A.4.c.iii	Agriculture/forestry/fishing: national fishing	Disaggregate to main fuel types		
1.A.5.a	Other, stationary (including military)			
1.A.5.b	Other, mobile (including military, land-based and recreational boats)			
1.B.1.a	Fugitive emission from solid fuels: coal mining and handling			
1.B.1.b	Fugitive emission from solid fuels: solid fuel transformation			
1.B.1.c	Other fugitive emissions from solid fuels			
1.B.2.a.i	Exploration production, transport			
1.B.2.a.iv	Refining/storage			
1.B.2.a.v	Distribution of oil products			
1.B.2.a.vi	Geothermal energy extraction			
1.B.2.b	Natural gas			
1.B.2.c	Venting and flaring	If this category is key, the inventory compiler should determine which subcategories are significant.		
2.A.1	Cement production			
2.A.2	Lime production			
2.A.3	Limestone and dolomite use			
2.A.4	Soda ash production and use			
2.A.5	Asphalt roofing			
2.A.6	Road paving with asphalt			
2.A.7.a	Quarrying and mining of minerals other than coal			
2.A.7.b	Construction and demolition			
2.A.7.c	Storage, handling and transport			
2.A.7.d	Other mineral products	If this category is key, the inventory compiler should determine which subcategories are significant.		

Source	e categories to be assessed in key category analysis	Special considerations		
Category code ^b	Category title ^b			
2.B.1	Ammonia production			
2.B.2	Nitric acid production			
2.B.3	Adipic acid production			
2.B.4	Carbide production			
2.B.5.a	Other chemical industry	If this category is key, the inventory compiler should determine which subcategories are significant.		
2.B.5.b	Storage, handling and transport	If this category is key, the inventory compiler should determine which subcategories are significant.		
2.C.1	Iron and steel production			
2.C.2	Ferroalloys production			
2.C.3	Aluminium production			
2.C.5.a	Copper production			
2.C.5.b	Lead production			
2.C.5.c	Nickel production			
2.C.5.d	Zinc production			
2.C.5.e	Other metal production	If this category is key, the inventory compiler should determine which subcategories are significant.		
2.C.5.f	Storage, handling and transport	If this category is key, the inventory compiler should determine which subcategories are significant.		
2.D.1	Pulp and paper			
2.D.2	Food and drink			
2.D.3	Wood processing			
2.E	Production of POPs (persistent organic pollutants)			
2.F	Consumption of POPs and heavy metals (e.g. electrical and scientific equipment)	If this category is key, the inventory compiler should determine which subcategories are significant.		
2.G Other production, consumption, storage, transportation or handling of bulk products		If this category is key, the inventory compiler should determine which subcategories are significant.		
3.A.1	Decorative coating application			

Source	e categories to be assessed in key category analysis	Special considerations		
Category code ^b	Category title ^b			
3.A.2	Industrial coating application			
3.A.3	Other coating application	If this category is key, the inventory compiler should determine which subcategories are significant.		
3.B.1	Degreasing			
3.B.2	Dry cleaning			
3.C	Chemical products			
3.D.1	Printing			
3.D.2	Domestic solvent use including fungicides			
3.D.3	Other product use			
4.B.1.a	Cattle dairy			
4.B.1.b	Cattle non-dairy			
4.B.2	Buffalo			
4.B.3	Sheep			
4.B.4	Goats			
4.B.6	Horses			
4.B.7	Mules and asses			
4.B.8	Swine			
4.B.9.a	Laying hens			
4.B.9.b	Broilers			
4.B.9.c	Turkeys			
4.B.9.d	Other poultry			
4.B.13	Other			
4.D.1	Synthetic N-fertilisers			
4.D.2.a	Farm-level agricultural operations including storage, handling and transport of agricultural products	If this category is key, the inventory compiler should determine which subcategories are significant.		
4.D.2.b	Off-farm storage, handling and transport of bulk agricultural products	If this category is key, the inventory compiler should determine which subcategories are significant.		
4.D.2.c	N-excretion on pasture range and paddock unspecified	If this category is key, the inventory compiler should determine which subcategories are significant.		
4.F	Field burning of agricultural wastes			

Source	e categories to be assessed in key category analysis	Special considerations		
Category code ^b	Category title ^b			
4.G	Agriculture Other	If this category is key, the inventory compiler should determine which subcategories are significant.		
6.A	Solid waste disposal on land			
6.B	Wastewater handling			
6.C.a	Clinical waste incineration			
6.C.b	Industrial waste incineration			
6.C.c	Municipal waste incineration			
6.C.d	Cremation			
6.C.e	Small-scale waste burning			
6.D	Other waste	If this category is key, the inventory compiler should determine which subcategories are significant.		
7.A	Other (included in national total for entire territory)			
^a In some cases ^b The categorie	s, inventory compilers may modify this list of categories to reflect parties s should include the respective codes and be consistent with the most re-	cular national circumstances. ecent NFR terminology.		

2.4 Methodological approaches to identify key categories

The methodology follows the IPCC approach to produce pollutant-specific key categories and covers Approaches 1 and 2 for both level and trend assessments. The two approaches developed by the IPCC, and described below, for performing key category analysis can be used to identify the key categories for each pollutant. Both approaches identify key categories in terms of their contribution to the absolute level of national emissions and to the trend of emissions.

- In Approach 1, key categories are identified using a predetermined cumulative emissions threshold. Key categories are those which, when summed together in descending order of magnitude, cumulatively add up to 80 % of the total level (³).
- In Approach 2, key categories can be derived by inventory compilers if category uncertainties or parameter uncertainties are available. Under Approach 2, categories are sorted according to their contribution to uncertainty.

Results of Approach 2 are additional to Approach 1. If Approach 1 and Approach 2 assessments have been performed, it is good practice to report the results of Approach 2 analysis in addition to the results of Approach 1. Results of both Approaches 1 and 2 should be used when setting priorities to inventory preparation.

 $[\]binom{3}{3}$ The predetermined threshold is based on an evaluation of several inventories and is aimed at establishing a general level where a significant percentage of inventory uncertainty will be covered by key categories. The final category that should be defined as *key* is that category for which the cumulative total is exactly equal to, or exceeds the 80 % threshold. This approach is consistent with that recommended by IPCC for the determination of key sources.

An additional step (see Appendix A) can be used for countries wanting to aggregate their pollutant-specific key category analysis to a single list based on weightings.

In addition to making a quantitative determination of key categories (or where quantitative assessments are not possible), it is good practice to consider qualitative criteria (for more detail, see subsection 2.4.3).

Figure 2-2 illustrates how inventory compilers can determine which approach to use for the identification of key categories.




Figure 2-2 Decision tree to identify key categories

2.4.1 Approach 1 to identify key categories

Approach 1 to identifying key categories assesses the influence of various categories of sources on the *level*, and, possibly, the *trend* of the national inventory. When the inventory estimates are available for several years, it is good practice to assess the contribution of each category to both the level and trend of the national inventory. If only a single year's inventory is available, a level assessment can only be performed.

Approach 1 can readily be accomplished using a spreadsheet analysis. Table 2-2 (level) and Table 2-3 (⁴) (trend) illustrate the format of the analysis. Separate calculation sheets (⁵) are suggested for each pollutant and for the level and trend assessments because it is necessary to sort the results of the analysis for each pollutant. It is more difficult to track the process if the analyses are combined in the same table.

Level assessment

The contribution of each source category to the total national inventory level is calculated according to equation (1) (level assessment (Approach 1)):

Key category level assessment = | source category estimate | / total contribution

$$L_{x,t} = E_{x,t} / \sum E_t \tag{1}$$

Where:

$\boldsymbol{L}_{x,t}$	=	level assessment for source x in latest inventory year (year t)				
$\boldsymbol{E}_{x,t}$	=	value of emission estimate of source category x in year t				
$\sum E_t$	=	total contribution, which is the sum of the emissions in year t , calculated using the aggregation level chosen by the country for key category analysis				

Key categories according to equation (1) are those that, when summed together in descending order of magnitude, add up to 80 % of the sum of all $L_{x,t}$. Table 2-2 presents a format that can be used for the level assessment.

 Table 2-2
 Spreadsheet for the Approach 1 analysis — level assessment

A^4	\mathbf{B}^4	C ⁴	D	Е	F
NFR category code	NFR category	Pollutant	Latest year estimate $E_{x,t}$	Level assessment $L_{x,t}$	Cumulative total of column E

^{(&}lt;sup>4</sup>) This guidance is copied from the 2006 IPCC Guidelines for National Greenhouse Gas Inventories and has been modified to provide column headings relevant for air pollutant emissions.

 $^(^{5})$ The 2006 IPCC Guidelines for National Greenhouse Gas Inventories provides an Excel spreadsheet tool for these calculations.

Total			$\sum E_t$	1		
-------	--	--	------------	---	--	--

Where:

column A:	code of NFR categories
column B:	description of NFR categories including fuel type
column C:	pollutant
column D:	value of emission estimate of category x in latest inventory year (year t)
column E:	level assessment following equation (1)
column F:	cumulative total of column F

Inputs to columns A–D will be available from the inventory. The total of column D presents the emissions. In column E, the level assessment is computed according to equation (1). Once the entries in column E are computed, the categories in the table should be sorted in descending order of magnitude according to column E. After this step, the cumulative total summed in column E can be calculated into column F. Key categories are those that, when summed together in descending order of magnitude, add up to 80 % of the total in column F. Where the method is applied correctly, the sum of entries in column E must be 1. The rationale for the choice of the 80 % threshold for the Approach 1 builds on Rypdal & Flugsrud (2001) and is also presented in the IPCC Guidelines, Chapter 7, subsection 7.2.1.1. This rationale suggests application of some pragmatic reduction to the threshold because of the number of pollutants for which emissions are estimated. Analysis of recent reporting shows that an 80 % cut-off captures a maximum of 75 % of categories for NMVOC reporting where emissions are important for a large number of categories.

The level assessment should be performed for the base year of the inventory (if applicable) and for the latest inventory year (year *t*). If estimates for the base year have changed or been recalculated, the base year analysis should be updated. Key category analysis can also be updated for other recalculated years. In many cases, however, it is sufficient to derive conclusions regarding methodological choice, resource prioritisation or QA/QC procedures without an updated key category analysis for the entire inventory time series. Any category that meets the threshold for the base year or the most recent year should be identified as *key*. However, the interpretation of the results of the key category analysis should take longer time series than the most recent year into account if key category analyses are available. If a category has been *key* for all or most previous years according to either the level or trend assessments or both (the two assessments should be considered separately), they should be identified as *key* in the latest year estimate except in cases where a clear explanation can be provided why a category may no longer be *key* in any future years.

Trend assessment

The purpose of the trend assessment is to identify categories that may not be large enough to be identified by the level assessment, but whose trend is significantly different from the trend of the overall inventory, and should therefore receive particular attention. The trend assessment can be calculated according to equation (2) if more than one year of inventory data are available.

$$T_{x,t} = \frac{E_{x,0}}{\sum E_0} \bullet \left[\left[\frac{\left(E_{x,t} - E_{x,0} \right)}{E_{x,0}} \right] - \frac{\left(\sum E_t - \sum E_0 \right)}{\sum E_0} \right]$$
(2)

Where:

 $T_{x,t}$ = trend assessment of source category x in year t as compared to the base year (year 0) or starting year of the inventory

$$E_{x,t}$$
 and $E_{x,0}$ = values of estimates of source category x in year t and 0 respectively
 $\sum E_t$ and $\sum E_0$ = total inventory estimates in years t and 0 respectively

The trend of a category refers to the change in the source category emissions over time; it is computed by subtracting the base year, or the starting year (year 0), estimate for source category x from the latest inventory year (year t) estimate and dividing by the value of the base year estimate.

The total trend refers to the change in the total inventory emissions over time, computed by subtracting the year 0 estimate for the total inventory from the latest year (year t) estimate and dividing by the value of the year 0 estimate.

In circumstances where the year 0 emissions for a given category are zero, the expression may be reformulated to avoid zero in the denominator (see equation (3)).

$$T_{x,t} = E_{x,t} / \sum E_0 \tag{3}$$

Where:

 $T_{x,t}$ = trend assessment of source category x in year t as compared to the base year (year 0) or starting year of the inventory

$$\sum E_0$$
 = total inventory estimates in years t and 0, respectively

$$E_{x,t}$$
 = values of estimates of source category x in years t

The trend assessment identifies categories whose trend is different from the trend of the total inventory, regardless of whether the category trend is increasing or decreasing, or is a sink or source. Categories whose trends diverge most from the total trend should be identified as *key*, when this difference is weighted by the level of emissions of the category in the base year.

Table 2-3 outlines a format that can be used for the Approach 1 trend assessment.

A^4	B ⁴	C ⁴	D	Е	F	G	Н
NFR category code	NFR category	Pollutant	Base year estimate $E_{x,0}$	Latest year estimate $E_{x,t}$	Trend assessment $T_{x,t}$	% Contribution to trend	Cumulative total of column G
Total					$\sum T_t$	1	

 Table 2-3
 Spreadsheet for the Approach 1 analysis — trend assessment

Where:

column A	:	code of NFR categories (⁴)
column B	:	description of NFR categories (⁴)
column C	:	pollutant (⁴)
column D	:	year 0 estimate of emissions from the national inventory data
column E	:	latest year estimate of emissions from the most recent national inventory data
column F	:	trend assessment from equation (2) (from equation (3) for zero base year emissions)
column G	:	percentage contribution of the category to the total of trend assessments in last row of column F, i.e. $T_{x,t}$ / $\sum T_t$
column H	:	cumulative total of column G, calculated after sorting the entries in descending order of magnitude according to column G

The entries in columns A, B, C and E should be identical to those used in Table 2-2, for the Approach 1 analysis — level assessment. The base year estimate in column D is always entered, while the latest year estimate in column E will depend on the year of analysis. The value of $T_{x,t}$ should be entered in column F for each category of sources, following equation (2), and the sum of all the entries entered in the total line of the table. The percentage contribution of each category to the total of column F should be computed and entered in column G. The categories (i.e. the rows of the table) should be sorted in descending order of magnitude, based on column G. The cumulative total of column G should then be computed in column H. Key categories are those that, when summed together in descending order of magnitude, add up to more than 80 % of the total of column F.

The trend assessment treats increasing and decreasing trends similarly. However, for the prioritisation of resources, there may be specific circumstances where countries may not want to invest additional resources in the estimation of key categories with decreasing trends. Underlying reasons why a category showing strong decreasing trends could include activity decrease, mitigation measures leading to reduced emission factors or abatement measures (e.g. chemical or metals production) changing the production processes.

In particular for a long-term decline of activities (not volatile economic trends) and when the category is not *key* from the level assessment, it is not always necessary to implement higher tier methods or to collect additional country-specific data if appropriate explanations can be provided why a category may not become more relevant again in the future. This could be the case, for example, for emissions of some pollutants (e.g. lead) from road vehicles in countries where switching to lead-free fuel reduced emissions significantly. Regardless of the method chosen or the importance of the category in particular years, countries should endeavour to use the same method for all years in a time series. Therefore, it may be more appropriate to continue using a higher tier method if it had been used in previous years.

Categories, identified using *key* in the trend assessment, which show declining trends, such as the introduction of abatement measures or other emission reduction measures, should be prioritised as appropriate estimation of these emissions will be important. Irrespective of the methodological choice, inventory compilers should clearly and precisely explain and document categories with strongly decreasing trends and should apply appropriate QA/QC procedures.

2.4.2 Approach 2 to identify key categories

Approach 2 is an enhancement on Approach 1 and helps to promote the importance of categories that have a high uncertainty. The approach is based on the results of the uncertainty analysis described in the Chapter 5, Uncertainties, in this Guidebook. Inventory compilers are encouraged to use Approach 2 in addition to Approach 1 because it will provide additional insight into the categories with high uncertainties and will assist in prioritising activities to improve inventory quality and reduce overall uncertainty.

Application of uncertainty estimates to identify key categories

Uncertainty estimates based on the uncertainty Approach 1 described in Chapter 5, Uncertainties, are sufficient for this purpose; however, estimates based on the uncertainty Approach 2 should be used when available. The category uncertainties are incorporated by weighting the Approach 1 level and trend assessment results according to the category percentage uncertainty. The key category equations are presented below.

Level assessment

Equation (4) describes the Approach 2 level assessment including uncertainty. This equation should be used to modify the equation in column E of Table 2-2 to add in the uncertainty component.

$$LU_{x,t} = \left(L_{x,t} \bullet U_{x,t}\right) / \sum \left[\left(L_t \bullet U_t\right)\right]$$
(4)

Where:

$LU_{x,t}$	=	level assessment for category x in latest inventory year (year t) with uncertainty
$\boldsymbol{L}_{x,t}$	=	computed as in equation (1)

 $U_{x,t}$ = category percentage uncertainty in year *t* calculated as described in Chapter 5, Uncertainties, and reported in column G of Table 6-1 in the same chapter. If the uncertainty is asymmetrical, the larger uncertainty

should be used. The relative uncertainty will always have a positive sign.

After computing level assessment with uncertainty, results should be sorted according to decreasing order of magnitude, as in Approach 1. The key categories are those that add up to 80 % of the sum of all $LU_{x,t}$. This 80 % was the basis for the derivation of the threshold used in the Approach 1 analysis. The categories identified by the level assessment with uncertainty that are different from categories identified by Approach 1 should then be highlighted and added to those also to be treated as additional key categories. In addition, the order of key categories identified by Approach 2 may be of use for those who are planning to improve inventories.

Trend assessment

Equation (5) shows how the Approach 2 Trend Assessment can be expanded to include uncertainty. This equation should be used to modify the equation in column F of Table 2-3 to add in the uncertainty component.

$$TU_{x,t} = \left(T_{x,t} \bullet U_{x,t}\right) \tag{5}$$

Where:

$TU_{x,t}$	=	trend assessment for category x in latest inventory year (year t) with uncertainty
$T_{x,t}$	=	trend assessment computed as in equation (2)
$U_{x,t}$	=	category percentage uncertainty in year t calculated as described in Chapter 5, Uncertainties. Note that this is the same uncertainty as in the total of column G of Table 6-1 in the same chapter and not the uncertainty assessment for trend. The relative uncertainty will always have a positive sign.

After computing trend assessment with uncertainty, results should be sorted according to decreasing order of magnitude. The key categories are those that add up to 80 % of the total value of the total $TU_{x,t}$. This 80 % was the basis for the derivation of the threshold used in the Approach 1 analysis. The key categories according to trend assessment with uncertainty should be treated as key categories and should be added to the list of key categories from Approach 1, if they are different from categories identified by Approach 1. In addition, the order of key categories identified by Approach 2 may be of use for those who are planning to improve inventories.

Incorporating Monte Carlo analysis

In Chapter 5, Uncertainties, Monte Carlo analysis is presented as the Approach 2 for quantitative uncertainty assessment. Whereas the Approach 1 uncertainty analysis is based on simplified assumptions to develop uncertainties for each category, Monte Carlo types of analysis can handle large uncertainties, complex probability density functions, correlations or complex emission estimation equations. The output of the Approach 2 uncertainty analysis can be used directly in equations (4) and (5). If uncertainties are asymmetrical, the larger percentage difference between the mean and the confidence limit should be used.

Monte Carlo analysis or other statistical tools can also be used to perform a sensitivity analysis to directly identify the principal factors contributing to the overall uncertainty. Thus, a Monte Carlo or similar analysis can be a valuable tool for a key category analysis. Inventory compilers are encouraged to use the method, for example, to analyse more disaggregated subcategories (by modelling correlations), emission factors and activity data separately (to identify key parameters rather than key categories). The use of these methods should be properly documented.

2.4.3 Qualitative criteria to identify key categories

In some cases, the results of Approach 1 or Approach 2 analysis of key categories may not identify all categories that should be prioritised in the inventory system. If quantitative key category analysis has not been carried out due to lack of completeness in the inventory, it is good practice to use qualitative criteria to identify key categories. The criteria below address specific circumstances that may not be readily reflected in the quantitative assessment. These criteria should be applied to categories not identified in the quantitative analysis. Those that meet the qualitative criteria should be added to the list of key categories. Although it is important to implement a trend assessment as part of good practice if data are available, it is particularly important to consider these criteria if the trend assessment has not been compiled.

- *Mitigation techniques and technologies*: if emissions from a category have decreased through the use of mitigation techniques, it is good practice to identify such categories as *key*. This will ensure that such categories are prioritised within the inventory and that better quality estimates are prepared to reflect the mitigation effects as closely as possible. It will also ensure that the methods used are transparent with respect to mitigation which is important for assessing inventory quality.
- *Expected growth*: the inventory compiler should assess which categories are likely to show an increase in emissions in the future. The inventory compiler may use expert judgement to make this determination. It is encouraged to identify such categories as *key*.
- No quantitative assessment of uncertainties performed: where Approach 2 including uncertainties in the key category analysis is not used, inventory compilers are still encouraged to identify categories that are assumed to contribute most to the overall uncertainty as *key*; the largest reductions in overall inventory uncertainty can be achieved by improving the estimates of categories having higher uncertainties. The qualitative consideration should take into account whether any methodological improvements could reduce uncertainties significantly. This could, for example, be applied to a small net flux results from the subtraction of large emissions, which can imply a very high uncertainty.
- *Completeness*: neither Approach 1 nor Approach 2 gives correct results if the inventory is not complete. The analysis can still be performed, but there may be key categories among those not estimated. In these cases it is good practice to examine qualitatively potential key categories that are not yet estimated quantitatively by applying the qualitative considerations above. The inventory of a country with similar national circumstances can also often give good indications of potential key categories. Chapter 3, Data collection, suggests methods to approximate activity data that can be used to compile preliminary estimates of emissions in a category. This preliminary analysis can thus be used to conclude whether a category potentially is *key* and prioritise data collection in that category.

The qualitative assessment should aim to introduce those categories within the quantitative ranking system, described in subsections 2.4.1 and 2.4.2, for each pollutant.

2.5 Reporting and documentation

It is good practice to clearly document the results of the key category analysis in an appropriate inventory report, such as the Informative Inventory Report (IIR), as specified in the UNECE EMEP Reporting Guidelines (UNECE, 2009). Documentation should include a series of tables by pollutant showing the prioritised categories, explaining the choice of method for each category (using Tier 1, 2 or 3 notations) and listing the criteria by which each category was identified as *key* (e.g. Approach 1, Approach 2, level or trend, or qualitative) using the following notations e.g. L1, L2, T1 or T2 where:

- L = key category according to level assessment
- T = key category according to trend assessment
- Q = key category according to qualitative criteria
- 1 = Approach 1 (basic cumulative on magnitude of emission)
- 2 = Approach 2 (applying uncertainty weightings)

Where categories are key and only Tier 1 methods are used a description of the difficulties in estimating using higher tiers should also be described.

2.6 Examples of key category analysis

Examples of the application of Approach 1 are shown in Table 2-4 and Table 2-5. Both the level and the trend assessment were conducted using estimates of emissions from the national inventory of Sweden (IVL, 2006). Currently this analysis does not show the recommended level of disaggregation by fuel. Although a qualitative assessment was not conducted in this example, it was not anticipated that additional categories would have been identified.

Examples of the results of the Approach 1 level assessment are shown in Table 2-4 with key categories in bold. Table 2-5 presents an Approach 1 trend key category analysis. Examples of Approach 2 level and trend assessments can be seen in the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (Chapter 4, Tables 4.9 and 4.10). Table 2-6 finally summarises the results of the key category analysis for NO_x for Sweden.

NOTE:

For air quality pollutants there should be a table, or tables, per pollutant as the emissions for different pollutants cannot be added together unless an optional approach for combining results of individual key category analyses is implemented (as specified in Appendix A). In addition, the examples below present the 80 % threshold suggested for key category analysis for air quality pollutants.

A^4	\mathbf{B}^4	C ⁴	D	E	F
A NFR	NFR category	Pollutant	Latest year estimate	Level	r Cumulative
category	itir K category	1 onutant	$E_{x,t}$	assessment	total of column
code			₽X,I	$L_{x,t}$	E
1.A.3.b.iii	Road transport:	NO _X	44.87	0.26	0.26
1110000	heavy-duty vehicles	I.O.A	1107	0.20	0.20
1.A.3.b.i	Road transport:	NO _X	28.12	0.16	0.42
	passenger cars	I.O.A	20112	0.10	
1.A.2.f	Other (specify in a	NO _X	21.29	0.12	0.54
1.1.1.2.1	covering note)	ΠOX		0.12	0.24
1.A.1.a	Public electricity and	NO _X	12.84	0.07	0.62
	heat production	I.O.A		0.07	0.02
2.D.1	Pulp and paper	NO _X	10.98	0.06	0.68
1.A.4.c.ii	Off-road vehicles and	NO _X	10.19	0.06	0.74
1010100	other machinery	I.O.A	10117	0.00	
1.A.3.d.ii	National navigation	NOx	5.96	0.03	0.77
1.A.2.d	Pulp, paper and print	NO _X	5.91	0.03	0.81
	- arp, paper and print	I.O.A	••• -		— see note
1.A.3.b.ii	Road transport: light-duty	NO _x	5.90	0.03	0.84
1.1.1.0.0.11	vehicles	I to x	0.90	0.05	0.01
1.A.4.c.iii	National fishing	NO _x	4.53	0.03	0.86
1.A.4.b.i	Residential plants	NO _x	3.98	0.02	0.89
1.A.3.e.ii	Other mobile sources and	NO _x	3.19	0.02	0.90
1.11.0.0.11	machinery	I to x	5.17	0.02	0.90
1.A.2.c	Chemicals	NO _x	2.28	0.01	0.92
1.A.3.a.ii.(ii)	Civil aviation (domestic,	NO _x	2.08	0.01	0.93
1.1.1.0	cruise)	I to x	2.00	0.01	0.95
1.A.4.b.ii	Household and gardening	NO _x	1.78	0.01	0.94
1.1.1.1.0.11	(mobile)	110 1	1., 0	0.01	0.9
1.A.1.b	Petroleum refining	NO _x	1.44	0.01	0.95
1.A.3.c	Railways	NO _x	1.36	0.01	0.95
1.A.5.b	Other, mobile (including	NO _x	1.24	0.01	0.96
	military)	1.0x	- · · · · ·	0.01	0.20
1.A.2.a	Iron and steel	NO _x	1.09	0.01	0.97
2.B.5	Other (specify in a	NO _x	1.05	0.01	0.97
	covering note)	1.0x		0.01	
2.C	Metal production	NO _x	0.95	0.01	0.98
2.A.7	Other including non-fuel	NO _x	0.68	0.00	0.98
	mining & construction	1.0x	0.00	0.00	0.20
	(specify in a covering				
	note)				
1.A.2.e	Food processing,	NO _x	0.62	0.00	0.99
	beverages and tobacco	1.0%	0.02	0.00	
1.A.3.a.ii.(i)	Civil aviation (domestic,	NO _x	0.45	0.00	0.99
w.m.(i)	LTO)	x	0.10	0.00	0.77

 Table 2-4. Example of Approach 1 level assessment (with key categories in bold)

A^4	\mathbf{B}^{4}	C ⁴	D	Е	F
NFR category	NFR category	Pollutant	Latest year estimate $E_{x,t}$	Level assessment	Cumulative total of column
code			Ayt	$L_{x,t}$	Ε
1.A.4.a	Commercial/institutional	NO _x	0.42	0.00	0.99
1.A.4.c.i	A/F/F Stationary	NO _x	0.35	0.00	0.99
2.B.2	Nitric acid production	NO _x	0.24	0.00	1.00
1.A.3.b.iv	Road transport: mopeds & motorcycles	NO _x	0.22	0.00	1.00
1.B.2.c	Venting and flaring	NO _x	0.16	0.00	1.00
1.A.1.c	Manufacture of solid fuels and other energy industries	NO _x	0.16	0.00	1.00
1.A.2.b	Non-ferrous metals	NO _x	0.09	0.00	1.00
6.C	Waste incineration	NO _x	0.09	0.00	1.00
1.B.1.c	Other (specify in a covering note)	NO _x	0.08	0.00	1.00
1.B.2.a.iv	Refining/storage	NO _x	0.02	0.00	1.00
6.C	Waste incineration	NO _x	0.09	0.00	1.00
1.B.1.c	Other (specify in a covering note)	NO _x	0.08	0.00	1.00
1.B.2.a.iv	Refining/storage	NO _x	0.02	0.00	1.00
1.B.2.a.vi	Other	NO _x	0.00	0.00	1.00

This example is based on the 2006 inventory of Sweden, where fuel type was not separated as recommended in these *Guidelines*. This does not affect categories identified as *key*. Note: Values in columns E and F are rounded. The final category that should be defined as *key* is that category for which the cumulative

Note: Values in columns E and F are rounded. The final category that should be defined as *key* is that category for which the cumulative total is exactly equal to, or exceeds the 80 % threshold. This approach is consistent with that recommended by the IPCC for the determination of key sources.

Α	В	С	D	Е	F	G	н
NFR category			$E_{x,0}$	$\boldsymbol{E}_{x,t}$	Trend assessment	% Contri-	Cumulative
code	NFR category	Pollutant	(Gg)	(Gg)	$T_{x,t}$	bution to trend	total of column G
1.A.3.b.i	Road transport: passenger cars	NO _x	105.58	28.12	0.10	0.43	0.43
1.A.3.b.iii	Road transport: heavy-duty vehicles	NO _x	58.78	44.87	0.04	0.17	0.60
2.D.1	Pulp and paper	NO _x	10.46	10.98	0.02	0.07	0.67
1.A.1.a	Public electricity and heat production	NO _x	14.44	12.84	0.02	0.07	0.74
1.A.2.f	Other (specify in a covering note)	NO _x	34.04	21.29	0.01	0.03	0.77
1.A.4.c.iii	National fishing	NO _x	4.51	4.53	0.01	0.03	0.80
1.A.3.d.ii	National navigation	NO _x	7.89	5.96	0.00	0.02	0.82
1.A.4.a	Commercial/institution	NO _x	3.32	0.42	0.00	0.02	0.84

 Table 2-5
 Example of Approach 1 trend assessment (with key categories in bold)

Α	В	С	D	Е	F	G	Н
			$\boldsymbol{E}_{x,0}$	$E_{x,t}$	Trend	%	
NFR category			$\mathbf{L}_{x,0}$	$\mathbf{L}_{x,t}$	assessment		Cumulative
code	NFR category	Pollutant	(Gg)	(Gg)	т	bution to	total of column G
			(Gg)	(Gg)	$T_{x,t}$	trend	commin G
	al						
1.A.5.b	Other, mobile	NO _x	4.63	1.24	0.00	0.02	0.86
1	(including military)	I to X			0.00	0.02	0.00
1.A.3.e.ii	Other mobile sources	NO _x	3.33	3.19	0.00	0.02	0.88
1.1.1.5.0.11	and machinery	I tox	5.55	5.17	0.00	0.02	0.00
1.A.2.c		NO _x	1.99	2.28	0.00	0.02	0.90
1 4 4 1 :	Chemicals Desidential plants	NO	9.22	2.00	0.00	0.02	0.01
1.A.4.b.i	Residential plants	NO _x		3.98	0.00	0.02	0.91
1.A.4.c.ii	Off-road vehicles and	NO _x	16.32	10.19	0.00	0.02	0.93
1 4 2 a ii (ii)	other machinery Civil aviation	NO	2.04	2.09	0.00	0.01	0.04
1.A.3.a.ii.(ii)		NO _x	2.04	2.08	0.00	0.01	0.94
1 4 2 4	(domestic, cruise)	NO	0.27	5.01	0.00	0.01	0.05
1.A.2.d	Pulp, paper and print	NO _x	9.37	5.91	0.00	0.01	0.95
1.A.4.b.ii	Household and	NO _x	2.33	1.78	0.00	0.01	0.96
	gardening (mobile)						
1.A.3.b.ii	Road transport:	NO _x	9.74	5.90	0.00	0.01	0.96
	light-duty vehicles						
2.B.2	Nitric acid production	NO _x	1.15	0.24	0.00	0.01	0.97
2.A.7	Other including	NO _x	0.73	0.68	0.00	0.00	0.97
	non-fuel mining &						
	construction						
1.A.2.e	Food processing,	NO _x	1.58	0.62	0.00	0.00	0.98
	beverages and tobacco						
1.A.1.b	Petroleum refining	NO _x	2.16	1.44	0.00	0.00	0.98
1.A.3.c	Railways	NO _x	2.09	1.36	0.00	0.00	0.98
1.A.3.b.iv	Road transport:	NO _x	0.09	0.22	0.00	0.00	0.99
	mopeds & motorcycles						
2.B.5	Other	NO _x	1.58	1.05	0.00	0.00	0.99
1.B.2.a.iv	Refining/storage	NO _x	0.27	0.02	0.00	0.00	0.99
1.A.2.a	Iron and steel	NO _x	1.74	1.09	0.00	0.00	0.99
1.B.1.c	Other (specify in a	NO _x	0.36	0.08	0.00	0.00	0.99
	covering note)						
1.B.2.c	Venting and flaring	NO _x	0.09	0.16	0.00	0.00	1.00
1.A.4.c.i	Stationary	NO _x	0.79	0.35	0.00	0.00	1.00
6.C	Waste incineration	NO _x	0.02	0.09	0.00	0.00	1.00
1.A.1.c	Manufacture of solid	NO _x	0.43	0.16	0.00	0.00	1.00
	fuels and other energy						
	industries						
1.A.3.a.ii. (i)	Civil aviation	NO _x	0.68	0.45	0.00	0.00	1.00
	(domestic, LTO)						
1.A.2.b	Non-ferrous metals	NO _x	0.20	0.09	0.00	0.00	1.00

Α	В	С	D	Ε	F	G	Н
NED optogory			$E_{x,0}$	$\boldsymbol{E}_{x,t}$	Trend assessment	% Contri-	Cumulative
NFR category code	NFR category	Pollutant	(Gg)	(Gg)	$T_{x,t}$	bution to trend	total of column G
1.B.2.a.vi	Other	NO _x	0.02	0.00	0.00	0.00	1.00
2.C	Metal production	NO _x	1.72	0.95	0.00	0.00	1.00
	s based on 2006 inventory of S ect categories identified as <i>key</i> .		fuel type wa	s not separate	ed as recommend	ded in these	e Guidelines.

Table 2-6 Summary of key category analysis (for NO_x)

Α	В	С	D	Е
NFR Category Code	NFR Category	Pollutant	Identification criteria	Comments ^a
1.A.3.b.iii	Road transport: heavy-duty vehicles	NO _x	L1,T1	1.A.3.b.iii
1.A.3.b.i	Road transport: passenger cars	NO _x	L1,T1	1.A.3.b.i
1.A.2.f	Other (specify in a covering note)	NO _x	L1,T1	1.A.2.f
1.A.1.a	Public electricity and heat production	NO _x	L1,T1	1.A.1.a
2.D.1	Pulp and paper	NO _x	L1,T1	2.D.1
1.A.4.c.ii	Off-road vehicles and other machinery	NO _x	L1	1.A.4.c.ii
1.A.3.d.ii	National navigation	NO _x	L1	1.A.3.d.ii
1.A.2.d	Pulp, paper and print	NO _X	L1	1.A.2.d
1.A.4.c.iii	National fishing	NO _x	T1	1.A.4.c.iii
^a Approach 1 use	ed only for NO _x .			

Quantitative method used: Approach 1

3 References

IPCC (2000) Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories, IPCC National Greenhouse Gas Inventories Programme, Institute for Global Environmental Strategies (IGES), Hayama, Japan (www.ipcc-nggip.iges.or.jp/public/gp/english/).

IPCC (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories, IPCC National Greenhouse Gas Inventories Programme, Global Environmental Strategies (IGES), Hayama, Japan (www.ipcc-nggip.iges.or.jp/public/2006gl/index.htm).

IVL (2006) Personal Communications, Kindbom, Karin.

Rypdal, K. & Flugsrud, K. (2001), 'Sensitivity Analysis as a Tool for Systematic Reductions in GHG Inventory Uncertainties', *Environmental Science and Policy*, Vol. 4 (2–3), pp. 117–135.

Statistics Finland (2005), *Greenhouse gas emissions in Finland 1990–2003*, National Inventory Report to the UNFCCC (United Nations Framework Convention on Climate Change), 27 May 2005.

UNECE (2009) Guidelines for reporting emission data under the Convention on Long-range Transboundary Air Pollution, ECE/EB.AIR/97.

4 Point of enquiry

Enquiries concerning this chapter should be directed to the co-chairs of the task force on emission inventories and projections. Please refer to the TFEIP website (<u>www.tfeip-secretariat.org/</u>) for the contact details of the current co-chairs.

Appendix A Combining air pollutant specific key categories

Once the trend and level key categories have been identified and ranked for each pollutant they can be combined. This is an optional procedure. This can be done by scoring each key category according to its position in the ranked pollutant key category list. The scoring applied can be tailored to national circumstances. A possible approach is to sum the percentage contributions across all pollutants to generate a single combined ranked list for each category.

This should be done for the level and the trend assessments using Approach 2, if uncertainty data are available, and Approach 1 if not.

Table A-1 shows the components for a simple Approach 1 level assessment example for CO and NO_x prior to aggregation. Similar tables should be compiled for all pollutants for which inclusion in the combined key category analysis is desired.

NFR category code	NFR category	Pollutant	year	Absolute value of latest year estimate	Level assessment $L_{x,t}$ (%)	Cumulative total of column F (%)	Key category flag
1.A.3.b.i	Road transport: passenger cars	СО	1171301	<i>E_{x,t}</i> 1171301	40	40	КС
1.A.4.b.i	Residential plants	СО	436585.7	436585.7	15	55	КС
1.A.2.f	Other, manufacturing industries and Construction	СО	407963.1	407963.1	14	69	КС
1.A.2.a	Iron and steel	СО	233561.2	233561.2	8	77	КС
2.C	Metal production	СО	99647.03	99647.03	3	80	КС
1.A.4.b.ii	Household and gardening (mobile)	СО	72701.24	72701.24	2	83	
1.A.3.b.iv	Road transport: mopeds & motorcycles	СО	71785.3	71785.3	2	85	
1.A.3.b.ii		СО	69050.87	69050.87	2	87	
1.A.1.a	Public electricity and heat	СО	67651.24	67651.24	2	90	

 Table A-1
 Simple example of approach 1 level assessment for CO and NO_x

NFR category code	NFR category	Pollutant	year	Absolute value of latest year estimate	Level assessment $L_{x,t}(\%)$	Cumulative total of column F (%)	Key category flag
				$ E_{x,t} $			
	production						
1.A.3.b.ii i	Road transport: heavy-duty vehicles	СО	55408.77	55408.77	2	92	
1.A.3.a.ii . (i)	Civil aviation (domestic, LTO)	СО	51834.52	51834.52	2	93	
2.B.5	Other, chemical industry	СО	28935.83	28935.83	1	94	
6.C	Waste incineration	СО	22526.49	22526.49	1	95	
1.A.4.c.ii	Off-road vehicles and other machinery (A,F,F)	СО	21252.68	21252.68	1	96	
2 A 4	Soda ash production and use	СО	20876	20876	1	97	
1.A.4.c.i	Stationary (A,F,F)	СО	15564.15	15564.15	1	97	
1.A.1.c	Manufacture of solid fuels and other energy industries	СО	11613.7	11613.7	0	98	
1.B.2.c	Venting and flaring (oil and gas)	СО	11581.24	11581.24	0	98	
1.A.3.a.i. (ii)	International aviation (cruise)	СО	0	0	0	98	
1.A.3.d.ii		СО	8214.236	8214.236	0	98	
1.A.3.a.i. (i)	International aviation (LTO)	СО	0	0	0	98	
1.A.5.b	Other, mobile (including military)	СО	7327.805	7327.805	0	98	

NFR category code	NFR category	Pollutant	year	Absolute value of latest year estimate	Level assessment $L_{x,t}(\%)$	Cumulative total of column F (%)	Key category flag
				$ E_{x,t} $			
5.B	Forest and grassland conversion	СО	6982.847	6982.847	0	99	
1.B.1.b	Solid fuel transformation	СО	6609.735	6609.735	0	99	
6.D	Other, waste	СО	6435.548	6435.548	0	99	
5.E	Other (not included in national total)	СО	0	0	0	99	
7	Other (included in national total)	СО	5874.342	5874.342	0	99	
1.A.4.a	Commercial/ institutional	СО	5408.98	5408.98	0	100	
1.A.1.b	Petroleum refining	СО	4726.058	4726.058	0	100	
1.A.3.c	Railways	СО	4487.554	4487.554	0	100	
1.A.3.e ii	Other mobile sources and machinery	СО	1645.713	1645.713	0	100	
1.A.3.a.ii . (ii)	Civil aviation (domestic, cruise)	СО	1359.588	1359.588	0	100	
1.A.2.b	Non-ferrous metals	СО	962.006	962.006	0	100	
1.B.2.a.i	Exploration, production, transport (oil)	СО	460.3594	460.3594	0	100	
1.A.1.a	Public electricity and heat production	NO _x	349625.7	349625.7	22	22	КС
1.A.3.b.ii i	Road transport: heavy-duty vehicles	NO _x	289605.2	289605.2	18	39	КС
1.A.3.b.i	Road transport: passenger cars	NO _x	245229.4	245229.4	15	55	КС
1.A.2.f	Other, manufacturing	NO _x	242722.7	242722.7	15	70	KC

NFR category code	NFR category	Pollutant	year	Absolute value of latest year estimate $ E_{x,t} $	Level assessment $L_{x,t}$ (%)	Cumulative total of column F (%)	Key category flag	
	industries and construction							
1.A.3.a.i. (ii)	International aviation (cruise)	NO _x	138290.8	138290.8	0	70	КС	
1.A.4.b.i	Residential plants	NO _x	114788.9	114788.9	7	77	КС	
1.A.1.c	Manufacture of solid fuels and other energy industries	NO _x	65729.83	65729.83	4	81		
1.A.3.d.ii		NO _x	63271.82	63271.82	4	85		
1.A.3.b.ii		NO _x	59155.09	59155.09	4	88		
1.A.4.c.ii	Off-road vehicles and other machinery (A,F,F)	NO _x	57548.33	57548.33	4	92		
1.A.1.b	Petroleum refining	NO _x	29347.65	29347.65	2	94		
1.A.2.a	Iron and steel	NO _x	20324.54	20324.54	1	95		
1.A.5.b	Other, mobile (including military)	NO _x	20165.12	20165.12	1	96		
1.A.3.c	Railways	NO _x	19231.36	19231.36	1	97		
1.A.4.a	Commercial/ Institutional	NO _x	19174.46	19174.46	1	98		
1.A.3.a.i. (i)	International aviation (LTO)	NO _x	11131.72	11131.72	0	98		
1.A.3.a.ii . (ii)	Civil aviation (domestic, cruise)	NO _x	6168.31	6168.31	0	99		
1.A.3.e.ii	Other mobile sources and machinery	NO _x	5625.079	5625.079	0	99		

NFR category code	NFR category	Pollutant	year	Absolute value of latest year estimate $ E_{x,t} $	Level assessment $L_{x,t}$ (%)	Cumulative total of column F (%)	Key category flag
1.B.2.c	Venting and flaring (oil and gas)	NO _x	2315.35	2315.35	0	99	
1.A.3.a.ii . (i)	Civil aviation (domestic, LTO)	NO _x	2279.565	2279.565	0	99	
6.C	Waste incineration	NO _x	1769.171	1769.171	0	100	
2.C	Metal production	NO _x	1559.267	1559.267	0	100	
1.A.3.b.i v	Road transport: mopeds & motorcycles	NO _x	1180.17	1180.17	0	100	
1.A.4.b.ii	Household and gardening (mobile)	NO _x	903.5456	903.5456	0	100	
2.B.2	Nitric acid Production	NO _x	736.3512	736.3512	0	100	
1.A.4.c.i	Stationary (A,F,F)	NO _x	725.5066	725.5066	0	100	
1.B.2.a.i	Exploration, production, transport (oil)	NO _x	523.5352	523.5352	0	100	
6.D	Other, waste	NO _x	421.2438	421.2438	0	100	
1.B.1.b	Solid fuel transformation	NO _x	364.7564	364.7564	0	100	
2.B.5	Other, chemical industry	NO _x	360.2754	360.2754	0	100	
5.E	Other	NO _x	212.0381	212.0381	0	100	
5.B	Forest and grassland conversion	NO _x	198.2986	198.2986	0	100	

The individual percentage contribution to totals for each pollutant (L) can be seen in column 6.

A summation of the percentage contributions for each of the identified key categories across the pollutants provides a method for ranking all pollutants considered. Table A-2 shows the simple example combining the CO and NO_x from Table A-1 above with additional pollutants to provide a single ranking across all pollutants.

Sector code	Sector description	%		ibutions to tegories (c	Sum of KC %	Rank				
couc	uescription	СО	1	NMVOC	1	PM ₁₀	1	SO	contributions	
1.A.1.a	Public electricity and heat production		1113		22	6	5	60	92	1
1.A.2.f	Other, manufacturing industries and construction	14		3	15	13	15	17	78	2
1.A.3.b.i	Road transport: passenger cars	40		7	15	4	7		73	3
1.A.4.b.i	Residential plants	15		4	7	17	15		57	4
1.A.3.b.iii	Road transport: heavy-duty vehicles			2	18	5	7		32	5
4.B.1.a	Dairy		25						25	6
4.B.1.b	Non-dairy		23						23	7
3.D	Other, solvent and other product use (including products containing HMs (heavy metals) and POPs)			22					22	8
1.A.3.b.ii	Road transport: light-duty vehicles					7	11		18	9
4.B.9	Poultry		11			6			18	10
2.A.7	Other, mineral products (including non-fuel mining & construction)					7	4		12	11
1.A.3.b.vi	Road transport: automobile tyre and brake wear					6	5		11	12
4.D.1	Direct soil emission		11						11	13
1.A.4.c.ii	Off-road vehicles and other machinery (A/F/F)					4	5		9	14
1.A.2.a	Iron and steel	8							8	15
2.D.2	Food and drink			8				ſ	8	16

 Table A-2
 Final key category ranking across all pollutants considered in analysis.

Sector	Sector	% 0		ibutions to	Sum of	Rank				
code	ode description		r -	tegories (c	KC %					
		CO	NH ₃	NMVOC	NO _x	\mathbf{PM}_{10}	PM _{2.5}	SO _x	contributions	
	production									
4.B.8	Swine		8						8	17
6.C	Waste incineration						5		5	18
1.B.2.a.i	Exploration, production, transport (oil)			5					5	19
1.A.3.b.v	Road Transport: Gasoline evaporation			3					3	20
1.B.2.a.iv	Refining, storage (oil)			3					3	21
1.B.2.a.v	Distribution of oil products			4					4	22
1.B.2.b	Natural gas			4					4	23
1.B.2.c	Venting and flaring (oil and gas)			3					3	24
2.B.5	Other, chemical industry			5					5	25
2.C	Metal production					4			4	26
3.B	Degreasing and dry Cleaning			3					3	27
3.C	Chemical products, manufacture and processing			1					1	28