

Chemicals

Emerging chemical risks in Europe — ‘PFAS’




It is currently not possible to perform in-depth environmental and health risk assessments of all chemical substances in use in Europe because of the great variety of chemicals and their diverse uses. New and legacy chemicals continue to be released into Europe’s environment, adding to the total chemical burden on Europe’s citizens and ecosystems. Early identification of emerging risks is one of the activities of the European Environment Agency (EEA). This briefing summarises the known and potential risks to human health and the environment in Europe posed by a group of very persistent chemicals, the per- and polyfluorinated alkyl substances (PFAS).

- Comprising more than 4 700 chemicals, per and polyfluorinated alkyl substances (PFAS) are a group of widely used, man-made chemicals that accumulate over time in humans and in the environment.
- National monitoring activities have detected PFAS in the environment across Europe. The production and use of PFAS in products has resulted in the contamination of drinking water supplies in several European countries. In some highly polluted areas, concentrations of perfluorooctanoic acid (PFOA) and perfluorosulfonic acid (PFOS) in drinking water were above the limit value for individual PFAS proposed in the 2018 recast of the EU Drinking Water Directive (EC, 2017).
- Human biomonitoring has detected a range of PFAS in the blood of European citizens. Though the levels for the most prevalent, studied and regulated PFAS, PFOA and PFOS are decreasing, levels of more ‘novel’ PFAS are increasing. In some areas, concentrations of PFOA and PFOS in the most exposed citizens were above proposed benchmark levels for adverse effects in humans.
- Due to the large number of PFAS chemicals, a substance-by-substance risk assessment and management approach is not adequate to efficiently prevent risk to the environment and human health from a single PFAS or mixtures of them.
- Taking precautionary risk management actions for groups of chemicals and promoting the use of chemicals that are ‘safe-and-circular-by-design’ could help to limit future pollution.

What are PFAS and what are they used for?

PFAS are a group of more than 4 700 man-made chemicals (OECD, 2018), the two most well-known of which are perfluorooctanoic acid (PFOA) and perfluorooctane sulfonic acid (PFOS) (Box 1). PFAS are used in a wide variety of consumer products and industrial applications because of their unique chemical and physical properties, including oil and water repellence, temperature and chemical resistance, and surfactant properties. PFAS have been used in firefighting foams, non-stick metal coatings for frying pans, paper food packaging, creams and cosmetics, textiles for furniture and outdoor clothing, paints and photography, chrome plating, pesticides and pharmaceuticals. Very limited information is available regarding which specific PFAS are used in which applications and at what levels in Europe.

Box 1: PFAS are a group of organic chemicals that contain a stable (unreactive) fluoro-carbon segment. Polyfluorinated PFAS contain both fluoro-carbon and hydro-carbon segments where the non-fluorinated part can degrade and ultimately form perfluorinated PFAS acids, such as PFOA and PFOS. While the long-chain PFAS accumulate in humans, animals and sediment/soil, the short-chain PFAS accumulate in the environment (German EPA, 2017, 2018) due to their persistency and high mobility in air and water. The [OECD](#) provides further information on groups of PFAS.




Why are PFAS a concern?

PFAS either are, or degrade to, persistent chemicals that accumulate in humans, animals and the environment. This adds to the total burden of chemicals to which people are exposed (Evans et al., 2016) and increases the risk of health impacts. Of the relatively few well-studied PFAS, most are considered moderately to highly toxic, particularly for children's development. Figure 1 summarises current knowledge of the health impacts of PFAS.

Environment and health

Figure 1. Effects of PFAS on human health



Sources: US National Toxicology Program, (2016); C8 Health Project Reports, (2012); WHO IARC, (2017); Barry et al., (2013); Fenton et al., (2009); and White et al., (2011).

People most at risk of adverse health impacts are those exposed to high levels of PFAS, and vulnerable population groups such as children and the elderly. Fewer studies have investigated effects on biota (Land et al., 2018). Throughout life, people and animals accumulate PFAS in their bodies. In 2018, the European Food Safety Authority (EFSA) re-evaluated the multiple lines of evidence of PFOA and PFOS toxicities, which resulted in significantly lower provisional 'safe' limits, known as the 'tolerable weekly intake' (TWI) (EFSA, 2018). The assessment concluded that a considerable proportion of the European population is expected to exceed the TWI due to intake of PFAS from food and drinking water.

Costs to society arising from PFAS exposure are high, with the annual health-related costs estimated to be EUR 52-84 billion across Europe in a recent study (Nordic Council of Ministers, 2019). The study notes that these costs are likely underestimated, as only a limited range of health

Environment and health

effects (high cholesterol, decreased immune system and cancer) linked to exposure to a few specific PFAS were included in the estimates. In addition, PFAS pollution also affects ecosystems and generates costs through the need for remediation of polluted soil and water. Such costs are currently difficult to assess since information on the number and scale of sites contaminated with PFAS in Europe and on how PFAS impact ecosystems is lacking.

What are the main sources of environmental PFAS pollution?




- Production and use of PFAS have been the main sources of PFAS contamination over time (Wang et al., 2014a, 2014b; Hu et al., 2016) for instance from fluoropolymer production installations and from the use of PFAS-containing firefighting foams (Figure 1). Other sources include PFAS produced and applied to textiles and paper and painting/printing facilities (Danish EPA, 2014). Less is known about potential releases of PFAS from other uses such as oil extraction and mining (Kissa, 2001), and the production of medical devices, pharmaceuticals and pesticides (Krafft and Riess, 2015).
- PFAS in consumer products, such as textiles, furniture, polishing and cleaning agents and creams, may contaminate dust and air, while food contact materials can contaminate food (Nordic Council of Ministers, 2019; Danish EPA, 2018). Drugs and medical devices may be other sources.
- Emissions to the environment occur via industrial waste water releases, as well as emissions to air from industrial production sites followed by deposition onto soil and water bodies. Industrial and urban waste water treatment plants are also a significant source of PFAS, via air, water and sludge (Hamid, et al., 2016; Eriksson et al., 2017).
- Reuse of contaminated sewage sludge as fertilisers has led to PFAS pollution of soil (Ghisi et al., 2019) and water in Austria, Germany, Switzerland and the US (Nordic Council of Ministers, 2019). The recycling of PFAS containing materials such as food contact materials and the formation of volatile fluorinated gases during waste incineration (Danish EPA, 2019) are other possible sources of PFAS pollution.

Where are PFAS found in Europe's environment?

PFAS are ubiquitous in the aquatic environment and organisms (Valsecchi et al., 2013) across Europe, and have been detected in air, soil, plants and biota (Houde et al., 2006). Areas around industrial production, manufacturing and application sites have been found to be particularly contaminated by PFAS. This has led to contaminated drinking water around factories in Belgium, Italy and the Netherlands, and around airports and military bases in Germany, Sweden and the United Kingdom (IPEN, 2018; Hu et al., 2016). The total number of sites potentially emitting PFAS is estimated to be in the order of 100 000 in Europe (Nordic Council of Ministers, 2019).

Environment and health

Generally, regulated PFAS have been substituted with other short-chain and polymeric PFAS. Regrettably, several of these 'novel' PFAS and their short chain degradation products are also persistent. In particular, short-chain PFAS accumulate in the environment and have been found to contaminate surface, ground- and drinking water (Eschauzier et al., 2012; Sun et al., 2016; Gebbink et al., 2017), and accumulate in plants (Ghisi et al., 2019), which may lead to increases in human dietary exposure.

Novel PFAS are increasingly detected (Xiao, 2017) in European surface waters. PFAS water pollution has been identified in countries across Europe, including Austria,  Denmark, France,  Germany, the Netherlands and  Sweden, as well as outside the EU. Several PFAS are sufficiently volatile to be considered long-range transboundary air pollutants, implying that emissions outside Europe are transported into Europe where they may accumulate in cold areas such as the Arctic (EEA, 2017). The well-known and regularly monitored PFAS (mainly perfluorinated acids) account only for a fraction of the chemical burden from PFAS present in human blood, the environment and wildlife (Koch et al., 2019).

While both well-known and novel PFAS have been detected in drinking water in non-EU countries (Xiao, 2017; Kaboré et al., 2018; Dauchy, 2019), at present there is little monitoring data available in the EU for drinking water. A case study by the World Health Organization (WHO) documents the story of PFAS contamination of the drinking water of 21 municipalities in the Veneto region of Italy. Industrial activity in the area had polluted both surface waters and ground water, as well as the drinking water of approximately 127 000 citizens (WHO, 2017). Monitoring conducted by the authorities of the Veneto Region found PFOS in 63-100 % of the locations sampled and PFOA in 100 % of the sites.

For comparison, the European Commission proposed a limit value of 0.1 µg/L for each individual PFAS in the 2018 recast of the EU Drinking Water Directive. This draft limit value was exceeded by a factor of 130 for PFOS and 66 for PFOA in samples taken in the Veneto Region.


PFOS and their derivatives are included as a priority hazardous substance under the EU Water Framework Directive (EU, 2013), with a much lower Environmental Quality Standard (AA-EQS) limit value of 0.65 ng/L (0.00065 µg/L) in inland surface waters and 0.13 ng/L in seawater. Member States are due to report on compliance with the PFOS EQS by 2021. Samples taken in 2013 in Northern Europe exceeded this EQS in 27 % of river sites and 94 % of Baltic Sea and Kattegat seawater (Nguyen et al., 2017).


What are the main routes of human exposure to PFAS?

The main exposure pathways for human and environmental exposures are shown in Figure 2. For the general population, PFAS sources include drinking water, food, consumer products and dust (EFSA, 2018). In food, fish species at the top of the food chain and shellfish are significant sources of PFAS exposure. Livestock raised on contaminated land can accumulate PFAS in their meat, milk and eggs (Ingelido et al., 2018; Numata et al., 2014). Direct exposure may also come via skin creams and cosmetics (Danish EPA, 2018; Schultes et al., 2018) or via air from sprays and dust from PFAS-coated textiles. There is little knowledge on uptake via skin and the lungs, which can be severely affected by PFAS (Nørgaard et al., 2010; Sørli et al., 2020). Consumer exposure may also occur via other routes such as via floor, wood, stone, and car polishing and cleaning products. Groups that may be exposed to high concentrations of PFAS include workers and people eating or drinking water and foods contaminated via PFAS treated food contact materials (Susmann et al., 2019). Though PFAS are used in drugs and medical equipment, there is little information on exposure via these routes.

Environment and health

Figure 2: Typical PFAS exposure pathways



PFAS are transferred in the womb from mother to child and unless exposure decreases with age, the PFAS body burden increases due to bioaccumulation (Koponen et al., 2018). Evidence of internal PFAS exposure in humans is available from several national human biomonitoring studies conducted inside and outside Europe. Men generally have higher PFAS body burdens and serum levels (Ingelido et al., 2018) because they excrete fewer PFAS. For the most regulated PFAS, such as PFOA and PFOS, consistent declines have been observed over the past 10-20 years in Europe (e.g. in Belgium, Denmark, Finland, Germany, Spain and Sweden). This decrease in levels in humans is likely to result from reduced exposure as a result of regulatory and non-regulatory action on consumer products, such as food contact materials (Susmann et al., 2019) and  textiles (Greenpeace, 2017).

Despite the decreases in long-chain PFAS levels in human blood, concentrations of PFOA and PFOS measured in human blood still exceed the EFSA benchmark dose levels (known as

Environment and health

BMDL5). This is particularly true for children and highly exposed sections of the European population (Buekers et al., 2018). The BMDL5 reflect the concentration in blood at which critical effects occur (cholesterol effects for adults and immune-toxicity for children) and are the basis for the provisional TWIs for PFOA and PFOS (EFSA, 2018).

The above mentioned human biomonitoring study in the Veneto Region investigated human exposure to PFOA and PFOS in the period 2015-2016 among 257 Italian residents of contaminated areas and 250 residents of background areas (Ingelido et al., 2018). The PFOA blood concentrations of residents of contaminated areas were 9-64 times higher than those of the background population. For PFOS, the levels were 1.4-1.6 times higher. Levels of PFOA in the highly exposed population were 0.2 to 26 times greater than the EFSA BMDL5, while for PFOS, the figure was 0.3-1.3 times. EU research projects, such as Human Biomonitoring for Europe (HBM4EU) (Box 2), are currently working to produce a representative picture of PFAS exposure for the European population, as well as investigating links between exposure and health effects.

Box 2: The Human Biomonitoring for Europe (HBM4EU) initiative is a 5-year EU Horizon 2020 research programme designed to translate human biomonitoring science into policy-relevant knowledge. A main task within the project is to generate representative chemical exposure data for the European population through harmonised human biomonitoring. PFAS is one of the 18 HBM4EU priority substance groups investigated by HBM4EU to better understand exposure and effects on health.

How can consumers avoid PFAS?

It is difficult for citizens to totally avoid exposure to PFAS. Using PFAS-free personal care products and cooking materials and avoiding direct contact with PFAS-containing products helps to reduce exposure. Decreased exposure to PFAS may be achieved by using consumer products from green labels and buying brands free from PFAS. General and specific guidance to consumers and business on how to find PFAS-free alternatives is provided by consumer organisations and some national institutions (see Danish EPA, German EPA and Swedish KEMI).

Decreased exposure to PFAS may be achieved by using consumer products from green labels and buying brands free from PFAS



© Engin_Akyurt / Pixabay

What is being done in the EU and globally?

Measures to reduce PFAS pollution are in place, mainly addressing well-known PFAS substances and their precursors. PFOS and PFOA are listed under Annex A of the Stockholm Convention on persistent organic pollutants (POPs), implying that parties to the Convention should 'eliminate the production and use' of the chemicals.

At EU level, PFOS is restricted under the EU POPs Regulation (EU, 2019). PFOA and its precursors are currently restricted under the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH) regulation (EU, 2006), including their presence in products made or

Environment and health

imported into the EU. This will soon be replaced by a new restriction under the POPs Regulation, which will have more limited derogations, following a decision taken at the Stockholm Convention.


A number of other PFAS are on the REACH list of Substances of Very High Concern (SVHCs). In June 2019, GenX (a short-chain PFAS substitute for PFOA in fluoropolymer production) was the first chemical added to the SVHC list on the basis of its persistent, mobile and toxic properties posing a threat to drinking water and the environment. Several PFAS are on the Community Rolling Action Plan for evaluation over the coming years. As mentioned above, PFOA and PFOS are priority hazardous substances under the Water Framework Directive (EC, 2017; EU, 2000).

Across Europe, several countries have been active in monitoring PFAS in environmental media as well as in humans and products. Some countries have set national limit values for water and soil (Denmark, Germany, the Netherlands and Sweden), for textiles (Norway) and for food contact materials (Denmark). Several EU Member States have set drinking water limits for specific PFAS and for groups of PFAS (Dauchy, 2019). In June 2019, Denmark announced a ban on PFAS-treated food contact materials, to enter into force in 2020.

Looking ahead

With more than 4 700 known PFAS, undertaking substance-by-substance risk assessments and comprehensive environmental monitoring to understand exposure would be an extremely lengthy and resource-intensive process. As a result, complementary and precautionary approaches to managing PFAS are being explored.

This includes the regulation of PFAS as a class, or as subgroups, based on toxicity or chemical similarities. The proposal to establish a new 'group limit' value for PFAS of 0.5 µg/L, in addition to limits for 16 individual PFAS of 0.1 µg/L in drinking water under the recast of the EU Drinking Water Directive is currently under consideration at EU level. Such measures can be supported by cost-effective and targeted monitoring of PFAS in the environment to provide early warning signals of pollution.

In June 2019, the  European Council of Ministers (EC, 2019) highlighted the widespread occurrence of PFAS in the environment, products and people, and called for an action plan to eliminate all non-essential uses of PFAS (Cousins et al., 2019).

The move towards zero pollution requires that product life cycles are made safer from the start (Warner and Ludwig, 2016), based on the concept of safe-and-circular-by-design (van der Waals et al., 2019). This approach offers opportunities to protect the health of Europe's citizens and environment at the same time as driving innovation for safer chemicals.

References


Kissa, E., 2001, 'Fluorinated Surfactants and Repellents: Second Edition, Revised and Expanded Surfactant Science Series. Volume 97, xiv + 616 pp, Marcel Dekker, New York. 2001, ISBN 0-8247-0472-X.', *Journal of the American Chemical Society* 123(36), pp. 8882-8882 (DOI: 10.1021/ja015260a).

Barry, V., et al., 2013, 'Perfluorooctanoic Acid (PFOA) Exposures and Incident Cancers among Adults Living Near a Chemical Plant', *Environmental Health Perspectives* 121(11-12), pp. 1313-1318 (DOI: 10.1289/ehp.1306615).


Buekers, J., et al., 2018, 'Development of Policy Relevant Human Biomonitoring Indicators for Chemical Exposure in the European Population', *International Journal of Environmental Research and Public Health* 15(10), p. 2085 (DOI: 10.3390/ijerph15102085).

C8 Health Project Reports, 2012, 'C8 Science Panel Website' accessed 2 December 2019.


Cousins, I. T., et al., 2019, 'The concept of essential use for determining when uses of PFASs can be phased out', *Environmental Science: Processes & Impacts* 21(11), pp. 1803-1815 (DOI: 10.1039/C9EM00163H).

Danish EPA, 2014,  Screeningsundersøgelse af udvalgte PFAS forbindelser som jord- og grundvandsforurening i forbindelse med punktkilder, Miljøprojekt No 1600.

Danish EPA, 2018, Risk assessment of fluorinated substances in cosmetic products, Survey of chemical substances in consumer products No 169, accessed 2 December 2019.

Danish EPA, 2019,  Belysning af destruktion af visse POP-stoffer på konventionelle affaldsforbrændingsanlæg til forbrænding af hovedsageligt ikkefarligt og forbrændingseget affald, No 2085, accessed 2 December 2019.

Dauchy, X., 2019, 'Per- and polyfluoroalkyl substances (PFASs) in drinking water: Current state of the science', *Current Opinion in Environmental Science & Health* 7, pp. 8-12 (DOI: 10.1016/j.coesh.2018.07.004).

EC, 2017, Proposal for a  DIRECTIVE OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL on the quality of water intended for human consumption (recast), COD No 0332, accessed 2 December 2019.

EC, 2019, Towards a Sustainable Chemicals Policy Strategy of the Union - Council conclusions (10713/19).

EEA, 2017, The Arctic Environment, European perspectives on a changing Arctic, Publication No 7, accessed 2 December 2019.

EFSA, 2018, Risk to human health related to the presence of perfluorooctane sulfonic acid and perfluorooctanoic acid in food, accessed 2 December 2019.

Environment and health

Eriksson, U., et al., 2017, 'Contribution of precursor compounds to the release of per- and polyfluoroalkyl substances (PFASs) from waste water treatment plants (WWTPs)', *Journal of Environmental Sciences (China)* 61, pp. 80-90 (DOI: 10.1016/j.jes.2017.05.004).

Eschauzier, C., et al., 2012, 'Impact of Treatment Processes on the Removal of Perfluoroalkyl Acids from the Drinking Water Production Chain', *Environ. Sci. Technol.* 46(3), pp. 1708-1715 (DOI: 10.1021/es201662b).

EU, 2000, Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy (327).

EU, 2006, Regulation (EC) No 1907/2006 of the European Parliament and of the Council of 18 December 2006 concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH), establishing a European Chemicals Agency, amending Directive 1999/45/EC and repealing Council Regulation (EEC) No 793/93 and Commission Regulation (EC) No 1488/94 as well as Council Directive 76/769/EEC and Commission Directives 91/155/EEC, 93/67/EEC, 93/105/EC and 2000/21/EC (OJ L).

EU, 2013, Directive 2013/39/EU of the European Parliament and of the Council of 12 August 2013 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy (Text with EEA relevance). (2013/39/EU).

EU, 2019, Regulation (EU) 2019/1021 of the European Parliament and of the Council of 20 June 2019 on persistent organic pollutants (Text with EEA relevance.) (OJ L).

Evans, R. M., et al., 2016, 'Should the scope of human mixture risk assessment span legislative/regulatory silos for chemicals?', *Science of The Total Environment* 543, pp. 757-764 (DOI: 10.1016/j.scitotenv.2015.10.162).


Fenton, S. E., et al., 2009, 'Analysis of PFOA in dosed CD-1 mice. Part 2. Disposition of PFOA in tissues and fluids from pregnant and lactating mice and their pups', *Reproductive Toxicology (Elmsford, N.Y.)* 27(3-4), pp. 365-372 (DOI: 10.1016/j.reprotox.2009.02.012).

Gebbink, W. A., et al., 2017, 'Presence of Emerging Per- and Polyfluoroalkyl Substances (PFASs) in River and Drinking Water near a Fluorochemical Production Plant in the Netherlands', *Environmental Science & Technology* 51(19), pp. 11057-11065 (DOI: 10.1021/acs.est.7b02488).

German EPA, 2017, **Protecting the sources of our drinking water from mobile chemicals**, Umweltbundesamt, accessed 2 December 2019.

German EPA, J., 2018, **'PFC-Planet: Chemikalien in der Umwelt'**, Umweltbundesamt, accessed 2 December 2019.

Ghisi, R., et al., 2019, 'Accumulation of perfluorinated alkyl substances (PFAS) in agricultural plants: A review', *Environmental Research* 169, pp. 326-341 (DOI: 10.1016/j.envres.2018.10.023).

Greenpeace, 2017,  **PFC Revolution in the Outdoor Sector**, accessed 2 December 2019.


Environment and health

Hamid, H. and Li, L., 2016, 'Role of wastewater treatment plant (WWTP) in environmental cycling of poly- and perfluoroalkyl (PFAS) compounds', *Ecocycles* 2(2) (DOI: 10.19040/ecocycles.v2i2.62).

Houde, M., et al., 2006, 'Biological Monitoring of Polyfluoroalkyl Substances: A Review', *Environmental Science & Technology* 40(11), pp. 3463-3473 (DOI: 10.1021/es052580b).

Hu, X. C., et al., 2016, 'Detection of Poly- and Perfluoroalkyl Substances (PFASs) in U.S. Drinking Water Linked to Industrial Sites, Military Fire Training Areas, and Wastewater Treatment Plants', *Environmental Science & Technology Letters* 3(10), pp. 344-350 (DOI: 10.1021/acs.estlett.6b00260).

Ingelido, A. M., et al., 2018, 'Biomonitoring of perfluorinated compounds in adults exposed to contaminated drinking water in the Veneto Region, Italy', *Environment International* 110(October 2017), pp. 149-159 (DOI: 10.1016/j.envint.2017.10.026).

IPEN, 2018,  **Fluorine-free firefighting foams (3F) viable alternatives to fluorinated aqueous film-forming foams (AFFF)**, Independent Expert Panel Convened by IPEN Stockholm Convention POPRC-14 Rome.

Kaboré, H. A., et al., 2018, 'Worldwide drinking water occurrence and levels of newly-identified perfluoroalkyl and polyfluoroalkyl substances', *Science of The Total Environment* 616-617, pp. 1089-1100 (DOI: 10.1016/j.scitotenv.2017.10.210).


Koch, A., et al., 2019, 'Towards a comprehensive analytical workflow for the chemical characterisation of organofluorine in consumer products and environmental samples', *TrAC Trends in Analytical Chemistry* (DOI: 10.1016/j.trac.2019.02.024).

Koponen, J., et al., 2018, 'Longitudinal trends of per- and polyfluoroalkyl substances in children's serum', *Environment International* 121, pp. 591-599 (DOI: 10.1016/j.envint.2018.09.006).

Krafft, M. P. and Riess, J. G., 2015, 'Per- and polyfluorinated substances (PFASs): Environmental challenges', *Current Opinion in Colloid & Interface Science* 20(3), pp. 192-212 (DOI: 10.1016/j.cocis.2015.07.004).

Land, M., et al., 2018, 'What is the effect of phasing out long-chain per- and polyfluoroalkyl substances on the concentrations of perfluoroalkyl acids and their precursors in the environment? A systematic review', *Environmental Evidence* 7(1), p. 4 (DOI: 10.1186/s13750-017-0114-y).

Nguyen, M. A., et al., 2017, 'Spatial distribution and source tracing of per- and polyfluoroalkyl substances (PFASs) in surface water in Northern Europe', *Environmental Pollution* 220, pp. 1438-1446 (DOI: 10.1016/j.envpol.2016.10.089).

Nordic Council of Ministers, 2019,  **The cost of inaction - A socioeconomic analysis of environmental and health impacts linked to exposure to PFAS**, TemaNord No 516.

Nørgaard, A. W., et al., 2010, 'Lung Damage in Mice after Inhalation of Nanofilm Spray Products: The Role of Perfluorination and Free Hydroxyl Groups', *Toxicological Sciences* 116(1), pp. 216-224 (DOI: 10.1093/toxsci/kfq094).

Environment and health

Numata, J., et al., 2014, 'Toxicokinetics of Seven Perfluoroalkyl Sulfonic and Carboxylic Acids in Pigs Fed a Contaminated Diet', *Journal of Agricultural and Food Chemistry* 62(28), pp. 6861-6870 (DOI: 10.1021/jf405827u).

OECD, 2018, 'About PFASs - OECD Portal on Per and Poly Fluorinated Chemicals', accessed 2 December 2019.

Schultes, L., et al., 2018, 'Per- and polyfluoroalkyl substances and fluorine mass balance in cosmetic products from the Swedish market: implications for environmental emissions and human exposure', *Environmental Science: Processes & Impacts* 20(12), pp. 1680-1690 (DOI: 10.1039/C8EM00368H).

Sørli, J. B., et al., 2020, 'Per- and polyfluoroalkyl substances (PFASs) modify lung surfactant function and pro-inflammatory responses in human bronchial epithelial cells', *Toxicology in vitro: an international journal published in association with BIBRA* 62, p. 104656 (DOI: 10.1016/j.tiv.2019.104656).

Sun, M., et al., 2016, 'Legacy and Emerging Perfluoroalkyl Substances Are Important Drinking Water Contaminants in the Cape Fear River Watershed of North Carolina', *Environmental Science & Technology Letters* 3(12), pp. 415-419 (DOI: 10.1021/acs.estlett.6b00398).

Susmann, H. P., et al., 2019, 'Dietary Habits Related to Food Packaging and Population Exposure to PFASs', *Environmental Health Perspectives* 127(10), p. 107003 (DOI: 10.1289/EHP4092).

US National Toxicology Program, 2016, **Toxicological Profile for Perfluoroalkyls**, accessed 2 December 2019.

Valsecchi, S., et al., 2013, 'Determination of perfluorinated compounds in aquatic organisms: a review', *Analytical and Bioanalytical Chemistry* 405(1), pp. 143-157 (DOI: 10.1007/s00216-012-6492-7).

van der Waals, J., et al., 2019, **Safe-by-design for materials and chemicals**, Zenodo, accessed 2 December 2019.


Wang, Z., et al., 2014a, 'Global emission inventories for C4-C14 perfluoroalkyl carboxylic acid (PFCA) homologues from 1951 to 2030, Part I: production and emissions from quantifiable sources', *Environment International* 70, pp. 62-75 (DOI: 10.1016/j.envint.2014.04.013).

Wang, Z., et al., 2014b, 'Global emission inventories for C4-C14 perfluoroalkyl carboxylic acid (PFCA) homologues from 1951 to 2030, part II: The remaining pieces of the puzzle', *Environment International* 69, pp. 166-176 (DOI: 10.1016/j.envint.2014.04.006).

Warner, J. C. and Ludwig, J. K., 2016, 'Rethink how chemical hazards are tested', *Nature News* 536(7616), p. 269 (DOI: 10.1038/536269a).

White, S. S., et al., 2011, 'Gestational and chronic low-dose PFOA exposures and mammary gland growth and differentiation in three generations of CD-1 mice', *Environmental Health Perspectives* 119(8), pp. 1070-1076 (DOI: 10.1289/ehp.1002741).

Environment and health

WHO, 2017,  Keeping our water clean: the case of water contamination in the Veneto Region, Italy, accessed 2 December 2019.

WHO IARC, 2017, Some Chemicals Used as Solvents and in Polymer Manufacture,.

Xiao, F., 2017, 'Emerging poly- and perfluoroalkyl substances in the aquatic environment: A review of current literature', Water Research 124, pp. 482-495 (DOI: 10.1016/j.watres.2017.07.024).

Identifiers

Briefing no. 12/2019

Title: **Emerging chemical risks in Europe — 'PFAS'**

PDF TH-AM-19-014-EN-N - ISBN 978-92-9480-196-8 - ISSN 2467-3196 -
doi: 10.2800/486213

HTML TH-AM-19-014-EN-Q - ISBN 978-92-9480-195-1 - ISSN 2467-3196 -
doi: 10.2800/02904

Published on 12 Dec 2019