Dutch bathing water quality in 2016







May 2017

Photo: © Peter Kristensen



BWD Report For the Bathing Season 2016 The Netherlands

The report gives a general overview of information acquired from the reported data, based on provisions of the Bathing Water Directive¹. The reporting process is described below, as well as state and trends of bathing water quality in the Netherlands.

1. BWD reporting in the season 2016

In 2016 bathing season, 718 bathing waters have been reported in the Netherlands. For each bathing water, five groups of parameters have been delivered²:

- identification data including name, location, geographic type of bathing water and availability to bathers;
- seasonal data including season start and end, national quality classification in present season, potential management measures and changes in quality;
- monitoring results disaggregated numerical values
 of two microbiological parameters intestinal
 enterococci and Escherichia coli (also known as E.
 coli), recorded at each water sample taken;
- abnormal situation periods periods of unexpected situations that have, or could reasonably be expected to have, an adverse impact on bathing water quality and on bathers' health; reporting is optional;

Bathing waters of the Netherlands in 2016						
Total reported	718					
Coastal	93					
Inland	625					
Max season period	153 days 1 May to 30 Sep					
Samples taken	6130					
Share of bathing waters with good or excellent water quality	91 %					
Reporting under Directive 2006/7/EC since	2009					

• *short-term pollution periods* – identifiable events that adversely affect water quality by faecal contamination; reporting is optional.

The authorities of the Netherlands report data according to the new BWD (2006/7/EC) since the season 2009. The data for the season 2016 were delivered to the European Commission by **20 December 2016**.

Altogether, **718 bathing waters** have been reported – 3.3% of all bathing waters in Europe. Out of all bathing waters in the Netherlands, 1.53% have been newly identified in 2016 season. 13% of bathing

¹ Directive BWD 2006/7/EC, available at http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=0]:L:2006:064:0037:0051:EN:PDF

² See the BWD Data Dictionary for detailed explanations: http://dd.eionet.europa.eu/datasets/latest/BWO_2006

waters in the Netherlands are of coastal type; the other 87% are inland. **6130 samples** were taken at bathing waters throughout the season – 9 per bathing water on average.

The maximum bathing season period was from 1 May to 30 September, i.e. 153 days altogether.

Detailed information on bathing waters is available from national portal at http://www.zwemwater.nl.

2. Assessment methodology³

During the bathing season, water samples are taken and analysed for two bacteria, *Escherichia coli* and intestinal enterococci which may indicate the presence of pollution, usually originating in sewage, livestock waste, bird faeces etc. The results of the analysis are used to assess the quality of the bathing waters concerned and to provide information to the public on the quality of water in the bathing sites concerned.

The monitoring requirements under the Directive are:

- taking a pre-season sample (taken shortly before the start of the bathing season) 4;
- a minimum of four samples per season⁵;
- a minimum of one sample per month⁶.

If these rules are satisfied, the bathing water is categorised as 'sampling frequency satisfied'. If not all monitoring requirements are fulfilled the bathing water is categorised as 'not enough samples'. 97.4% of bathing waters met the described monitoring requirements set by the Directive, while the rest did not satisfy monitoring requirements for different reasons: being new; having changed environmental conditions that might affect water quality classification; closed; not monitored due to legal issues, physical inaccessibility to the site etc.

 $^{^{3}}$ The methodology used by the EC and the EEA is described here, while results of assessment by national authorities may differ in individual cases.

⁴ A pre-season sample is taken into account at total number of samples per season.

⁵ Three samples are sufficient if the season does not exceed eight weeks or the region is subject to special geographical constraints.

⁶ If, for any reason, it is not possible to take the sample at the scheduled date, a delay of four extra days is allowed. Thus, the interval between two samples should not exceed 31 + 4 days.

Table 1 shows the statistics of bathing waters according to monitoring requirements.

Table 1: Bathing waters in 2016 according to compliance with BWD monitoring provisions

	Count	Share of total [%]	
BWs with sampling frequency satisfied (and are not new, are not subject			
to changes or were not closed in 2016)		97.4%	
These bathing waters have been monitored according to provisions and	699		
have complete dataset from the last assessment period. They have been			
quality-classified (excellent, good, sufficient, poor).			
BWs with sampling frequency not satisfied (and are not new, are not		0.0%	
subject to changes or were not closed in 2016)			
These bathing waters exist throughout the last assessment period but have	0		
not been monitored throughout the period according to provisions for	0		
various individual reasons. They may be quality-classified if there is an			
adequate volume of samples available for credible classification.			
BWs that are new, subject to changes or closed in 2016		2.6%	
These bathing waters do not have complete dataset for the last assessment			
period because they are new, have been subject to changes (that are likely	19		
to affect the classification of the bathing water) or have been closed. They			
cannot be quality-classified.			
Total number of bathing waters in 2016	718	100%	

Bathing waters where sampling frequency was not satisfied can still be quality assessed if at least four samples per season (three samples if the season does not exceed eight weeks or the region is subject to special geographical constraints) are available and equally distributed throughout the season. Assessment of bathing water quality is possible when the bathing water sample dataset is available for four consecutive seasons. Bathing waters are accordingly classified to one of the bathing water quality classes (excellent, good, sufficient, or poor).

The classification is based on pre-defined percentile values for microbiological enumerations, limiting the classes given in Annex I of the Directive. The Directive defines different limit values for coastal and inland waters.

Quality assessment is not possible for all bathing waters. In these cases, they are instead classified as either:

- not enough samples⁷;
- new8:
- changes9;
- closed¹⁰.

⁷ Not enough samples have been provided throughout the last assessment period (the last four bathing seasons or, when applicable, the period specified in Article 4.2 or 4.4).

⁸ Classification not yet possible because bathing water is newly identified and a complete set of samples is not yet available.

⁹ Classification is not yet possible after changes that are likely to affect the classification of the bathing water.

¹⁰ Bathing water is closed temporarily or throughout the bathing season.

3. Bathing water quality

The results of the bathing water quality in the Netherlands throughout the past period are presented in Figure 1 (for coastal bathing waters) and Figure 2 (for inland bathing waters). The previous reports are available on the European Commission's bathing water quality website¹¹ and the European Environment Agency's bathing water website¹².

3.1 Coastal bathing waters

In the Netherlands, 96.8% of all existing coastal bathing waters met at least sufficient water quality standards in 2016. See Appendix 1 for numeric data.




Figure 1: Coastal bathing water quality trend in the Netherlands. Note: the "At least sufficient" class also includes bathing waters of "Excellent" quality class, the sum of shares is therefore not 100%.

¹¹ http://ec.europa.eu/environment/water/water-bathing/index_en.html

¹² http://www.eea.europa.eu/themes/water/status-and-monitoring/state-of-bathing-water

3.2 Inland bathing waters

94.4% of all existing inland bathing waters were of at least sufficient water quality in 2016. See Appendix 1 for numeric data.




Figure 2: Inland bathing water quality trend in the Netherlands. Note: the "At least sufficient" class also includes bathing waters of "Excellent" quality class, the sum of shares is therefore not 100%.

4. Information regarding management and other issues

Bathing water information can be accessed through the "Zwemwater" portal available at http://www.zwemwater.nl. Bathing water quality and bathing water locations can be viewed using the map viewer. Besides the map viewer, the portal also provides information regarding safe bathing, bathing prohibitions, health risks, bathing water profiles etc.

At the bathing water sites large light blue information boards are present. The boards provide general information about the bathing area: a map showing the swimming area, bathing water quality, information about available facilities and potential prohibitions. In the case when the bathing water does not meet required quality standards, a special plate is put under the general information board.

5. Bathing water quality assessment presentation in online viewers

The European bathing water legislation focuses on sound management of bathing waters, greater public participation and improved information dissemination. More on the bathing and other water legislation can be found on the European Commission's website: http://ec.europa.eu/environment/water/index en.htm.

The bathing water section of the Water Information System for Europe (WISE) which is accessible at the EEA bathing water website (http://www.eea.europa.eu/themes/water/interactive/bathing/state-of-bathing-waters) allows users to view the bathing water quality at more than 21 000 coastal and inland sites across Europe. The WISE bathing water quality data viewer combines text and graphical visualisation, providing a quick overview of the bathing water's locations and achieved quality. Having access to bathing water information, citizens are encouraged to make full use of it and participate with their comments.

Appendix 1: Results of bathing water quality in the Netherlands from 2013 to 2016


Table 2: Bathing waters in the season 2016 according to quality

		Total number of bathing waters	of Excellent qua		At least sufficient quality		Poor quality		Quality classification not possible: not enough samples /new bathing waters/bathing waters subject to changes/closed	
			No	%	No	%	No	%	No	%
	2013	90	66	73.3	87	96.7	3	3.3	0	0.0
Coastal	2014	91	62	68.1	87	95.6	3	3.3	1	1.1
	2015	92	69	75.0	89	96.7	1	1.1	2	2.2
	2016	93	75	80.6	90	96.8	0	0.0	3	3.2
	2013	621	423	68.1	567	91.3	33	5.3	21	3.4
Inland	2014	624	441	70.7	577	92.5	32	5.1	15	2.4
l la	2015	622	461	74.1	580	93.2	23	3.7	19	3.1
	2016	625	462	73.9	590	94.4	19	3.0	16	2.6
	2013	711	489	68.8	654	92.0	36	5.1	21	3.0
Total	2014	715	503	70.3	664	92.9	35	4.9	16	2.2
	2015	714	530	74.2	669	93.7	24	3.4	21	2.9
	2016	718	537	74.8	680	94.7	19	2.6	19	2.6

Note: the class "At least sufficient" also includes bathing waters which are of excellent quality, the sum of shares is therefore not 100%.

Appendix 2: Bathing water quality map

Map 1: Bathing waters reported during the 2016 bathing season in the Netherlands



Source: National boundaries: EEA; Large rivers and lakes: EEA, WFD Article 3; Bathing waters data and coordinates: Dutch authorities; Digital Elevation Model over Europe (EU-DEM): EEA.